

ANTIMICROBIAL STEWARDSHIP IN AUSTRALIAN LIVESTOCK INDUSTRIES



© 2018 Australian Eggs Limited.

All rights reserved.

ISBN 978-1-920835-20-0

Antimicrobial Stewardship in Australian Livestock Industries

The views expressed, and the conclusions reached in this publication are those of the author and those that contributed and not necessarily those of persons consulted. Australian Eggs Limited shall not be responsible in any way whatsoever to any person who relies in whole or in part on the contents of this report.

This publication is copyright, however, Australian Eggs Limited encourages wide dissemination of its research, providing that it is clearly acknowledged. For any other enquiries concerning reproduction, contact the Sustainability Program Manager on 02 9409 6999.

Author Contact Details

Name: Dr. Kylie Hewson

Phone: 0422 760 736

Email: kylie@sativus.com.au

In submitting this report, the researcher has agreed to Australian Eggs Limited publishing this material in its edited form.

Australian Eggs Limited Contact Details:

Australian Eggs Limited

A.B.N: 6610 2859 585

Address: Suite 6.02, Level 6, 132 Arthur St
North Sydney NSW 2060

Phone: 02 9409 6999

Fax: 02 9954 3133

Email: research@australianeggs.org

Website: www.australianeggs.org.au

Published in November 2018

CONTENT

Foreword	4
Abbreviations	5
Introduction	6
Historical Australian Government AMS Efforts	8
The Australian Veterinary Association	11
Overview of the Contributing Industries	11
The 5R's for Australian Livestock AMS	13
The Australian Pork Industry	17
The Australian Poultry Industries	21
The Australian Red Meat Industries	29
The Australian Dairy Industry	34
Conclusion	37
Acknowledgements	37
References	38
Appendix 1	41
Appendix 2	43
Appendix 3	44
Appendix 4	46

FOREWORD

The following report on Antimicrobial Stewardship (AMS) within Australian livestock industries is the result of a collaborative effort, which has been coordinated by Australian Eggs on behalf of, and with approval from, other industries and relevant AMS experts.

This report provides an overview of historical and current practises relevant to AMS in each of the contributing industries and is primarily intended for the stakeholders who are interested to know how livestock industries operate in Australia. Although some industries have not contributed to this report, it does not in any way imply a lack of dedication from those industries to AMS.

The volume of information supplied for each often reflects the intensity of production rather than their level of commitment to AMS. To an extent, the amount of information provided reflects the varying efforts required to manage diseases in different livestock systems. An important long-term goal is that this report will be updated to include contributions from other Australian livestock industries.

INDUSTRY CONTACTS

For enquiries on further information related to antimicrobial stewardship in the industries included in this report, please refer to the relevant contact below.

<i>Industry</i>	<i>Organisation</i>	<i>Contact Email</i>
Chicken meat	Australian Chicken Meat Federation	acmf@chicken.org.au
Dairy	Dairy Australia	enquiries@dairyaustralia.com.au
Duck meat	Australian Duck Meat Association	g.parkinson@iinet.net.au
Egg	Australian Eggs	contacts@australianeggs.org.au
Grain-fed beef / Grass-fed beef / Sheep-meat	Meat & Livestock Australia	info@mla.com.au
Pork	Australian Pork Limited	research@australianpork.com.au
Poultry primary breeders	Specialised Breeders Australia	sbaadmin@specialisedbreeders.com.au
Turkey	Australasian Turkey Federation	info@turkeyfed.com.au

Advisory Contacts

<i>Organisation</i>	<i>Contact Email</i>
Australian Veterinary Association	melanie.latter@ava.com.au
Advanced Veterinary Therapeutics	swp@advet.com.au
NSW Department of Primary Industries	david.jordan@dpi.nsw.gov.au

ABBREVIATIONS

ACMF	Australian Chicken Meat Federation
ACV	Australian Cattle Veterinarians
ALFA	Australian Lot Feeder's Association
AMR	Antimicrobial Resistance
AMS	Antimicrobial Stewardship
APL	Australian Pork Limited
APV	Australian Pig Veterinarians
APVMA	Australian Pesticides and Veterinary Medicines Authority
ARC	Australian Research Council
AVA	Australian Veterinary Association
AVPA	Australasian Veterinary Poultry Association
DAWR	Australian Government Department of Agriculture and Water Resources
ESA	Egg Standards of Australia
GVP	Gross Value of Production
IDF	International Dairy Federation
JETACAR	Joint Expert Technical Advisory Committee on Antibiotic Resistance
LPA	Livestock Production Assurance
MHU	Minor Human Use
MLA	Meat & Livestock Australia
NHMRC	National Health and Medical Research Council
NHU	No Human Use
QA	Quality Assurance
RDC	Research and Development Corporation
RD&E	Research, Development and Extension

INTRODUCTION

“Antimicrobial resistance (AMR) is one of the biggest threats to human and animal health today” is the opening statement of the Australian Government website devoted to AMR (<https://www.amr.gov.au/>). This statement highlights the increasing national and international attention being given to the harmful consequences of antimicrobial resistance that may influence the health and welfare of current and future generations of animals and people. The Australian Government Department of Health and Department of Agriculture, now the Department of Agriculture and Water Resources, released “Responding to the Threat of Antimicrobial Resistance”, in 2015 which was Australia’s first National Antimicrobial Resistance Strategy covering the 2015-2019 period.

The overall goal of the National AMR Strategy is to minimise the development and spread of AMR to ensure the continued availability of effective antimicrobials for use in humans and animals (Australian Government, 2015). The National AMR Strategy describes seven objectives to successfully accomplish this goal, including the pivotal importance of implementing antimicrobial stewardship (AMS) practises across human health and animal care settings, to ensure appropriate and judicious prescription and administration of antimicrobials. This current document presents an overview of AMS initiatives already practiced across a variety of livestock operations in Australia.

What is Antimicrobial Stewardship?

‘Stewardship’ describes a situation where an individual takes responsibility for the care and management of something not owned by that individual. In the past, stewardship has applied to responsible care of forests, water, oceans, the environment in general (Darden, 1988) and to animal welfare. At the beginning of the 21st Century, the critical importance of the care and management of antimicrobials as precious, non-renewable resources has received heightened focus. The term ‘stewardship’ was applied to practises that minimise the need for antimicrobial use, as well as practises related to their use.

One of the most distinct descriptions of AMS was provided by two eminent researchers who are experts in appropriate antimicrobial use in animals, Luca Guardabassi from Denmark and John Prescott from Canada (Guardabassi and Prescott, 2015). They define AMS as, “the multifaceted and dynamic approaches

required to sustain the clinical efficacy of antimicrobials by optimizing drug use, choice, dosing, duration, and route of administration, while minimizing the emergence of resistance and other adverse effects”. “Multifaceted and dynamic” indicates that AMS is complex, involves many elements, requires relevant knowledge, and reflects the fact that its practise is not static. The prevention of disease in animals and optimisation of drug use, choice, dosing, duration and route of administration when disease occurs is situation specific and each circumstance requires a tailored approach.

Before the development and publication of Australia’s first National AMR Strategy, various sections of the Australian animal health community, particularly those devoted to livestock, had been refining production systems to minimise disease challenge to animals. This was achieved through a collaborative relationship between farmers and veterinarians, and their efforts encompassed many of the elements of what is now described as antimicrobial stewardship (AMS). The AMS elements have always been at the core of operations for livestock industries and their supporting veterinarians, due at least in part to the economic benefits of growing livestock that are free from disease. In many cases, the cost of treating or preventing disease with antimicrobials, and the resulting loss in productivity of a sick animal, outweighs the cost of improving biosecurity, hygiene practises and overall husbandry, which are key principles of good AMS. Therefore, no specific date can be ascribed as the ‘start’ of AMS in any livestock industry, as in many industries these principles have long been included and formalised as requirements under various quality assurance (QA) and accreditation programs.



The 5R framework for AMS

The 5R framework for AMS was developed to provide a systematic and comprehensive approach to AMS planning, implementation and monitoring, to allow a potentially complex process to be both practical and effective (Lloyd and Page, 2018; Page et al., 2014; Prescott and Boerlin, 2016; Scott Weese et al., 2013).

The AMS framework (Figure 1) includes five (5) essential components:

- Responsibility
- Review
- Reduce
- Refine
- Replace

which are described in detail in the relevant sections of this report.

‘Good Stewardship Practice’ describes the development, implementation and continual improvement of the AMS plan, the collaborative process between those responsible for the livestock and those responsible for supporting the health and welfare of the livestock. This includes tailoring approaches at the farm, business and enterprise levels.

The baseline and progress

The progressive introduction of biosecurity and infection prevention and control measures (including vaccination) in Australian livestock has significantly reduced the incidence of infections that require treatment. This proactive management, coupled with the limited number of antimicrobial agents available in Australia to treat bacterial disease in livestock, has resulted in the low incidence of AMR bacteria recovered from Australian pigs (Abraham et al., 2017; Kidsley et al., 2018; Obeng et al., 2014), meat chickens (Obeng et al., 2012a; Obeng et al., 2012b, 2014), laying chickens and their eggs (Obeng et al., 2014; Pande et al., 2015) and cattle (Barlow et al., 2015, 2017). The progressive introduction of AMS programs provides a solid foundation for maintaining and improving the current baseline of low antimicrobial use, low incidence of antimicrobial resistance and high animal health and welfare.

Compared to most other countries, Australia has an enviable reputation for low use of antimicrobial agents and low frequency of AMR. However, there are only ad hoc systems in place to capture this information, although there are efforts through the National AMR Strategy to dramatically improve availability of data (Australian Government, 2015).



Figure 1. The 5R framework for good antimicrobial stewardship.

HISTORICAL AUSTRALIAN GOVERNMENT AMS EFFORTS

In 1997, Australia became one of the first nations to embark on a comprehensive process of reform aimed at protecting humans and animals from the harmful effects of AMR infections. The work of the Joint Expert Technical Advisory Committee on Antibiotic Resistance (JETACAR), which produced the 'JETACAR report' in 1999 (JETACAR, 1999) set the scene for future efforts, and pointed out the significance of managing AMR in both humans and animals as a single system. This 'One Health' concept encourages "healthy people, healthy animals and a healthy planet". In 2012, a renewed focus arose from a heightened collaboration between the Australian Government's medical and veterinary authorities, with improvements aimed toward aligning with the One Health approach. As a result, Australia's First National Antimicrobial Resistance Strategy was released in 2015 (Australian Government, 2015), which was followed by an Implementation Plan in 2016 (Australian Government, 2016). The National AMR Strategy recognises that, to successfully combat AMR, the community needs a broad-based effort on AMS to be simultaneously implemented across all sectors in the animal health and human healthcare systems. The National AMR Strategy supports collaboration to build on previous efforts and implement new initiatives in the reduction of inappropriate antibiotic usage contributing to AMR in humans and animals. The Australian livestock industries and State/Territory and Commonwealth Governments are active participants in the implementation of this strategy.

The Australian public sector has historically implemented a suite of actions for supporting AMS in food animals. As Australia is a major global exporter of high-quality food products, governments and industry have a strong working partnership to manage all issues of food integrity, including the use of antimicrobials in animals. A cornerstone of AMS involves the work of the Australian Pesticides and Veterinary Medicines Agency (APVMA); a national agency that takes on responsibility for registration of chemicals (including antimicrobials) for use in animals (APVMA, 2018). The APVMA strictly applies a process of scientific review for new antimicrobial products which includes an antimicrobial resistance risk assessment to determine the possible impact of use on the health of Australians and sets risk management controls on use accordingly. This approach is uniquely conservative by

global standards; wherein there are no antimicrobial products registered for use in food animals in Australia that contain fluoroquinolones, carbapenems, colistin or fourth generation cephalosporins and the use of third generation cephalosporins in food animals is highly constrained. A comprehensive description of the APVMA's history and role in helping to coordinate Australia's human health and veterinary agencies to control AMR was released in 2017 (APVMA, 2017).

Complementing the Commonwealth regulations are those directed at 'control of use' of antimicrobials enacted by the states and territories of Australia. These regulations govern all aspects of the supply and use of antimicrobials, including compliance with APVMA imposed restrictions. Jurisdictions also implement their own legislation aimed at maintaining high professional standards in animal health by ensuring only veterinarians with current registration can prescribe scheduled substances, such as antimicrobials. Prescription of antimicrobials by a veterinarian has been progressively implemented since the 1970s, with additional antimicrobials made prescription-only as risk assessments were undertaken over time. Historic AMS initiatives in Australia since the 1950s are outlined in Figure 2. The commitment of the Commonwealth and State/Territory Governments, along with animal industries, is demonstrated through the work of Animal Health Australia, which coordinates industry-government initiatives aimed at improving animal health, welfare and biosecurity. The Government provides significant contributions to the annual 'Animal Health in Australia' report, which is a comprehensive summary of Australia's animal health status and system (AHA, 2018).

The Commonwealth and State/Territory governments are also active in supporting the Australian Veterinary Association's (AVA) efforts toward the implementation of good AMS principles by Australian veterinarians and through various mechanisms, encourage adoption of the "Guidelines for Prescribing, Authorising and Dispensing Veterinary Medicines 2005 (updated in 2013)", which is overseen by the AVA (AVA, 2013). The combined effect of Commonwealth and state/territory legislation ensures that antibiotics can only be administered to livestock when prescribed by a veterinarian and must be used according to label directions (APVMA, 2015). The use of antimicrobials, like all agricultural and veterinary chemicals, is controlled to ensure that residues do not exceed maximum residue limits in edible tissues of livestock at the time of slaughter. This requirement is monitored by the Australian Government's National Residue Survey, which was established by the Australian Government in the early 1960s (Australian Government, 1961).


Further, the Australian Strategic and Technical Advisory Group on AMR (ASTAG) was established in 2014 and is co-chaired by the Australian Chief Medical Officer and the Australian Chief Veterinary Officer, with membership from a broad range of relevant stakeholder groups. ASTAG is part of the broader AMR framework in Australia and researches and provides expert advice on AMR related issues, including the development and maintenance of an Australian-specific importance rating for antibacterial agents (ARSC, 2014; ASTAG, 2018).

Australian Commonwealth and State/Territory governments sponsor a broad range of research and development initiatives aimed at improving the stewardship of antimicrobials in the food animal sector. A major contribution is through financial support to the Rural Research and Development Corporations (RDCs; <http://www.ruralrdc.com.au/>) servicing individual industries (details provided elsewhere in this document). Research related to AMS is also sponsored through research granting institutions, such as the National Health and Medical Research Council (NHMRC) and the Australian Research Council (ARC). The Australian Government has directly supported several research projects, including AMR surveillance in food animals (Australian Government, 2007; Shaban et al., 2014), and a series of additional proof-of-concept surveillance projects between 2016-2018 (ACMF, 2018a; other reports not available at the time of this publication). They have also supported the development of an education program on AMS for veterinary graduates for Australian and NZ veterinary schools that is expected to be launched in late 2018/early 2019.

Commonwealth and State/Territory Governments continue to identify and prioritise opportunities for supporting and initiating activities related to improving veterinary AMS in Australia, with a renewed focus on collaborative One Health efforts.



Australian Veterinary Antimicrobial Stewardship Timeline



Compiled by Stephen Page. Designed by Ed Hewson, TKOAH.

Figure 2. Key historical veterinary antimicrobial stewardship actions that have progressed veterinary antimicrobial stewardship in Australia. Source: Dr. Stephen Page; references supplied in Appendix 1.

THE AUSTRALIAN VETERINARY ASSOCIATION

For more than 30 years, the AVA has been actively involved in fighting the emergence of AMR through the development of factsheets and guidelines for prescribing, authorising and dispensing veterinary medicines, as well as designing a code of practice to specifically support the judicious use of antimicrobials (all available, plus other resources from: <https://www.ava.com.au/amr>). Fighting AMR has been one of the AVA's three strategic priorities since 2016 (AVA, 2018).

The AVA is an active participant in Commonwealth One Health initiatives and the implementation of the National AMR Strategy (Australian Government, 2015), and there are several aspects of the AVA's program to address AMR. The AVA's policies on the use of veterinary medicines and documents related to the management of livestock health and disease prevention can be accessed at: <https://www.ava.com.au/about-us/policy-and-positions-1>.

Antibiotic prescribing guidelines

In 2017, the AVA embarked on a project in partnership with Animal Medicines Australia (AMA) to develop best-practice, evidence-based guidelines for the prescription of antibiotics to pigs, through funding provided by the Australian Government and Australian Pork Limited. Following completion of the pig guidelines, the AVA and AMA will jointly fund and facilitate the development of guidelines for the prescription of antibiotics to poultry, sheep, horses, and cattle (beef, dairy and feedlot).

Antimicrobial stewardship


Evidence-based guidelines for the prescription of antibiotics already exist for domesticated dogs and cats (Holloway, 2013) and in 2016, the AVA initiated the pilot trial of an AMS program for companion animal practices in Canberra. This trial was instigated by two Canberra animal health practitioners and was supported by a small working group of staff and volunteers. Results are expected to be published in 2019.

Community awareness

The AVA has participated in Antibiotic Awareness Week since 2012 and has been represented on the national organising committee of this event since 2013. Their involvement will continue as a way of helping to increase understanding of antibiotic use and resistance in animals among animal owners and human health professionals.

One Health policy agenda

AVA contributes to the ASTAG and is part of the monitoring and implementation of the National AMR Strategy. The implementation plan includes several key projects that are managed by the AVA (Australian Government, 2016).



OVERVIEW OF THE CONTRIBUTING INDUSTRIES

This report provides an overview of historical and current practises relevant to AMS in each of the contributing industries and is primarily intended for the stakeholders who are interested to know how livestock industries operate in Australia. Although some industries have not contributed to this report, it does not in any way imply a lack of dedication from those industries to AMS. Further, the volume of information supplied for each often reflects the intensity of production rather than their

level of commitment to AMS. To an extent, the amount of information provided reflects the varying efforts required to manage diseases in different livestock systems.

A summary of key industry information is provided in Table 1. All details in this section have been obtained from a wide-range of resources from the relevant industry organisations.

Table 1: Key Annual statistics for the contributing industries.

Industry	GVP (\$billion) ¹	Livestock Slaughtered (million) ²	Industry Representative Organisation	Industry RD&E Organisation	RD&E budget (\$million) ³	Contact
Chicken meat	2.60	652	Australian Chicken Meat Federation	AgriFutures Australia	4.0	chicken.org.au
Dairy	4.1	1.56 ⁴	Australian Dairy Farmers	Dairy Australia	61	dairyaustralia.com.au
Duck	0.12	10	Australian Duck Meat Association	<i>In development</i>	N/A	N/A
Egg	0.82	27.9 ⁵	Egg Farmers of Australia	Australian Eggs	4.5	eggfarmersaustralia.org
Pork	1.34	5.16	Australian Pork Limited	Australian Pork Limited / High Integrity Pork CRC	10.5 / 20	australianpork.com.au
Turkey	0.03	4.5	Australasian Turkey Federation	N/A	N/A	turkeyfed.com.au
Beef (feedlot and pasture)	12.14	23.3	Australian Lot Feeders Association / Cattle Council	Meat & Livestock Australia	104	mla.com.au
Sheep meat	3.56	70.2	Sheep Producers Australia			sheepproducers.com.au

¹ABS, June 2017 Value of Agricultural commodities produced www.abs.gov.au/ausstats/abs@.nsf/mf/7503.0, estimates made of proportion of poultry slaughters;

²ABARES Agricultural commodities: March quarter 2018 report 16-17 estimate; ³2016/17 annual reports - value of levied RD&E investment and does not include the value of RD&E undertaken within individual companies; ⁴Dairy herd; ⁵Pullets and layers; Australian Eggs Annual Report, 2017.

THE 5R'S FOR AUSTRALIAN LIVESTOCK AMS

This report highlights key efforts by each industry that contributed against each of the headings for the 5 R framework – Responsibility, Reduce, Refine, Replace, Review. This report is not designed to be comprehensive, but rather to provide avenues for obtaining further information as required by the reader of this report.

Responsibility

A successful AMS plan requires a shared responsibility between the livestock producer, who is responsible for maintaining animals to high standards of health and welfare and following all directions for use and implementing associated management changes, and the prescribing veterinarian, who accepts responsibility for the decision to use an antimicrobial agent. Responsibility must also engender and promote the enthusiastic support from all levels of each organisation, including senior management, veterinarians and those working in practices, on farms and own animals (Cunha, 2018).

If producers believe that there may be a health issue on their site, they are responsible for alerting their company (if contractually obligated), their veterinarian or relevant flock/herd health management professional, and/or the relevant State Government Authority (in the case of a suspected emergency animal disease). The Model Codes of Practice for the Welfare of Animals has specific codes for each species (available from CSIRO publishing <https://www.publish.csiro.au>). There are provisions stating that those people responsible for livestock must take reasonable actions to ensure the welfare of the animals under their care, which includes the responsibility to understand and follow vaccination, chemical and medication treatment instructions. There are also several other obligations of producers, such as the development of livestock health plans. These Model Codes of Practice are in the process of being turned in to 'Standards and Guidelines', with 'Standards' eventually being incorporated into relevant legislation (further information available at: <http://www.animalwelfarestandards.net.au/>).

There is also responsibility for those that have relevant knowledge of appropriate AMS to improve the knowledge of others through various methods, such as provision of codes of practices, best practice manuals and training. The AVA offers several further development and training

opportunities for Australian veterinarians (further information available at: <https://www.vetvoice.com.au/ec/veterinary-careers/continuing-professional-development>).

Reduce

Wherever possible, means of reducing the use of antimicrobials should be implemented. Biosecurity measures underpin animal health and welfare and are supported by meticulous hygiene, precision nutrition, vaccination, and expert animal husbandry. The combination of these measures will ensure infectious disease incidence (and need for antimicrobials) is minimised. High order biosecurity measures and the continuing improvement of and use of vaccines provide enormous support for low use of antimicrobials (Hoelzer et al., 2018a, b).

Although reducing the use of antimicrobials is just one aspect of AMS, reducing their use is a key driver for all livestock industries. In many cases, the cost of treating or preventing disease with antimicrobials and the resulting loss in productivity of sick animals, outweighs the cost of improving biosecurity, hygiene practises and overall husbandry.

Industry specific biosecurity manuals have been developed in consultation with the Australian Government to outline the recommended minimum biosecurity practices that should be employed to prevent incursion of pathogens into the herd or flock (biosecurity manuals available at: <http://www.farmbiosecurity.com.au/livestock>). These biosecurity manuals require ratification by members of Animal Health Australia, which include state and Commonwealth Government representatives and representatives from many animal industries. Several, if not all, of these biosecurity recommendations are captured as requirements through various QA programs, accreditation programs and farm contracts (where farmers are contracted to supply livestock).

A greater understanding of the physiological, nutritional and behavioural requirements of all livestock species over time, coupled with the implementation of tighter biosecurity controls, has led to improvements in the prevention of disease incursion in Australian livestock, which subsequently minimises the use of antimicrobials.

Optimal genetic variation of stock is achieved through controlled breeding programs, which include consideration of the robustness and resilience of livestock to disease challenges simultaneously with quality production traits. Significant advances in assisted breeding technologies have resulted in livestock that are more resistant to illness than previous generations, and improvements in this area are continually progressing (Colditz and Hine, 2016; Hermesch and Dominik, 2014).

As there is no current standard measure for the quantity of antimicrobials that can be used for benchmarking, it is difficult to ascertain the extent of reduction over time resulting from the implementation of new technologies and practices. Through various legislative and commercial requirements, veterinarians (and in the instances of contract farming; the company) are required to maintain records of antimicrobial prescription and/or use. However, the completeness of this data and the accessibility and ability to consolidate it is widely variable, as are the potential commercial-in-confidence considerations related to the data.

Refine

'Refinement' is defined as the right diagnosis and, if antimicrobials are required, the right drug, at the right time, at the right dose, through the right route, and for the right length of time. During the initial and ongoing reviews of an AMS plan, there are many elements of refinement that require attention. For example, improved detection of disease may allow earlier identification of animals for treatment, which can result in disease containment, improved animal health and potentially less ability for AMR to develop (Neethirajan, 2017; Richeson et al., 2018).

There are a limited number of antimicrobial agents with antibacterial activity that are approved for use in Australia. The agents approved for use in cattle, sheep, pigs and meat and egg laying poultry are presented in Appendix 2. Many approved agents are not used in human medicine and are not considered to present a risk of selecting AMR of significance to human health. There are very few agents approved for use in animals that are considered of importance to human health. All agents are appropriate candidates for a refinement strategy, and consideration of what constitutes appropriate use is of vital importance (Smith et al., 2018).

Australian livestock industries use diagnostic assays to detect and support confirmation of diseases in livestock. These assays are essential for evaluating whether an animal, herd or flock is suffering from viral, parasitic or bacterial diseases, and to appropriately direct

antimicrobial use (i.e. not using an antibiotic to treat a viral infection). There are assays available to determine the resistance of bacteria, but they are expensive and, as such, are not often used by veterinarians to determine resistance profiles prior to prescribing or administering antimicrobials.

The growth-promoting effect of antimicrobials can arise when an antimicrobial agent is used at sub-therapeutic doses as a feed supplement in food animals. While the mode of action is not fully understood, it is theorised that by suppressing commensal bacteria (which divert nutrition from the animal), more nutrients are able to be utilised by the animal for growth/production and less energy is required for maintaining the integrity of the gastrointestinal tract (Gaskins et al., 2002). Australian livestock industries have agreed to formalise a long-standing practice of not using antimicrobials of importance to human health for animal growth promotion purposes by progressing with the voluntary removal of growth promotion label claims from these antimicrobials (refer to the Office of the Australia Chief Veterinary Officer for more information). Antimicrobial manufacturers have removed (or are in the process of removing) all label claims for growth promotion from antimicrobials important for human health.

Application of antimicrobials is a key area for refinement of use and in most cases are administered via feed or drinking water. Antimicrobials are often administered through feed to allow mass medication of large numbers of animals and to achieve uniform distribution of medication throughout the herd/flock. It is also suitable for long-term medication, such as coccidiosis control. Medicating via feed also removes the risk of error and omission when dispersal is dependent on a farm worker administering medication in the water on a frequent (once or twice daily) basis. Medicated feed does have its limitations; it cannot be implemented quickly enough to treat rapidly spreading, or fast onset diseases. Implementation and cessation of medication can also be difficult to manage to avoid cross contamination when medicated and unmedicated feed may be stored sequentially in the same silo.

Water medication is also very useful for administering antimicrobials to large groups of animals, and it allows for better refinement of the treatment than feed medication because the decision to medicate can be made and implemented quickly and terminated more rapidly. A disadvantage of water medication is that sick animals often reduce or stop water intake, therefore, there is the risk that sick animals do not receive adequate medication for successful treatment. Not all antimicrobials are able to be administered via the water.

The prescribing guidelines being developed by the AVA will support livestock industries to further identify areas of possible refinement of antimicrobial use for disease prevention and treatment.

Replace

There is substantial national and global interest in finding non-antimicrobial approaches to support the continual health and welfare of animals. It is important that any measures taken to replace antimicrobials are based on sufficient evidence of effectiveness and safety in order to avoid unintended adverse consequences. Economic viability is an important consideration in livestock production where small profit margins are common. A significant focus in Australian livestock is on the development of vaccines to reduce disease incidence, as vaccination and improved hygiene and biosecurity is more cost effective than managing disease, disease-related productivity declines and welfare issues. Apart from vaccines, other alternatives to antimicrobials that are the subject of active investigation are enzymes (Hassan et al., 2018), phytochemicals (Lillehoj et al., 2018), microbial products (Seal et al., 2018), various immunoglobulins and host defence peptides (van Dijk et al., 2018) and products containing essential oils and feed additives containing short and medium chain fatty acids. These alternative products, including those that aren't classified as medications, are regularly trialled on-farm to determine efficacy as a replacement for the prevention or treatment of livestock diseases. The extent of this type of trialling is dependent on the individual producer/company and is often not publicly reported.

Vaccines are the primary focus for livestock industries for limiting the impact of animal disease and are therefore central to livestock AMS. Vaccines can be 'live', 'inactivated' or 'attenuated', and may also be from strains of bacteria that are from the farm itself ('autogenous') that are then supplied back to animals on that farm only. Australian livestock often require vaccines specific to strains of pathogen present in Australia (due, in part, to Australia's geographic isolation). There is a relatively small market for those vaccines in Australia, and an inability to cost-effectively sell small volumes of vaccines (for smaller producers to access). Australia's strict quarantine requirements, designed in part to protect Australian livestock from diseases prevalent overseas, also limit the supply of vaccines to the Australian livestock sectors. Although RD&E funding organisations are continually investing in efforts to develop additional vaccines, it can be cost-prohibitive to commercialise vaccines that are effective in Australia only.

Industries work with the regulatory authorities to help reduce the time between development of a vaccine candidate and registration of a product available for use in industry, and help to gain access to vaccines that may be available overseas. The PUBCRIS (Public Chemical Registration Information System; <https://portal.apvma.gov.au/pubcris>) database can be searched for current and historical information, including registration dates for vaccines available for Australian livestock. A comprehensive list of vaccines available is presented in Appendix 3. Autogenous vaccines may also be produced and administered under permit to prevent known local diseases (details available at <https://portal.apvma.gov.au/permits>). Vaccines are widely used where possible and recent technological advances may mean that more diseases can be controlled by vaccines in the future.

Review

'Review' involves recording and regularly reviewing the effectiveness of AMS practises, including the quantity and quality of antimicrobial use when required, and using a selection of the most appropriate metrics or descriptors to ensure that antimicrobial use practises reflects best practice (Collineau et al., 2017; EMA and ESVAC, 2017; Harbarth and Hackett, 2018; Le Maréchal et al., 2018; Versporten et al., 2018). In addition, an appropriate AMR surveillance program provides important guidance on progress and outcomes, and the relevance of AMS plan objectives (Jee et al., 2018; Simjee et al., 2018).

The AMS review process can take several different forms and includes the continual review and acknowledgement of improvements made through industry QA and other third-party QA accreditation programs, particularly components that relate to health, hygiene and husbandry. All livestock produced in Australia have State legislative requirements, as well as various requirements for QA depending on consumer and market requirements and, in most instances, there are industry-wide voluntary QA programs. All these approaches to QA and auditing require that accurate and detailed records are kept, including any medications administered to the livestock and production details; this provides numerous opportunities for review of antimicrobial use practises. There are currently no nationally adopted metrics for measuring quantity or quality of antimicrobial use in Australian livestock, and previous information available has been based on antimicrobial sales data (APVMA, 2014).

Surveillance for AMR in commensal bacteria supports AMS by providing information on the effectiveness, or otherwise, of various interventions. Several surveillance studies have been undertaken for various industries since the early 2000s and continue to be repeated, though usually with slight alterations as methodologies and economies for AMR testing improve. Some of these studies cover several livestock species (Australian Government, 2007), Australian pigs (Abraham et al., 2017; Kidsley et al., 2018; Obeng et al., 2014), meat chickens (Obeng et al., 2012a; Obeng et al., 2012b, 2014), laying chickens and their eggs (Obeng et al., 2014; Pande et al., 2015) and cattle (Barlow et al., 2015, 2017), with reports from recent projects in the pork, chicken meat and egg industries expected in 2018/19 and the red meat industries in 2019/20.



Australian Eggs



THE
AUSTRALIAN
PORK
INDUSTRY

THE AUSTRALIAN PORK INDUSTRY

The size and geographic location of Australian pig farms vary due to a range of factors, including climate, the availability and price of feed, environmental permits/licences, the demand and type of pork imports, currency fluctuations, and competition from other meat products. As of 2018, the Australian pork industry consists of approximately 400 producers that supply 90% of the total domestic production. A conservative estimate of pig holders outside of the commercial system is 1500-2000.

The size of pig farms vary from small, peri-urban operations to large, vertically integrated operations that can house tens of thousands of pigs on multiple farm sites linked by an overseeing company. Pigs may be reared indoors in conventional penned housing, or in deep-litter sheds. Outdoor pig units are where the animals are raised in paddocks or pens, operate in smaller areas and generally supply niche markets. Semi-intensive piggeries keep dry sows (those with no piglets and not lactating) outside, and the mating and growing portions of the piggery are kept inside.

About 90% of the Australian sow herd is produced under the Australian Pork Industry Quality Assurance Program (APIQ✓®). To comply with APIQ✓®, each farm must have a herd health program and a consultant pig veterinarian that oversees all medical treatments.

Live pigs and any pig genetic material (e.g. semen for breeding purposes) are not allowed to be imported into Australia, however, approximately 300,000 tonnes of pork is imported annually from several countries, which has steadily increased as amended quarantine arrangements allow for imports to be sourced from more international suppliers. Under current arrangements, all imported pork must be cooked in Quarantine Approved Premises regulated by DAWR to comply with requirements of the Pork Import Risk Assessment and manufactured into processed meat products. Up to 70% of the ham, bacon and smallgoods produced in Australia are made from imported pork.

Responsibility

The antimicrobial stewards for the Australian pork industry are the private and corporate veterinarians employed by, or working with, Australian pig producers. These veterinarians are required to be registered, and many are also members of the AVA and the Australian Pig Veterinarians (APV), a special interest group of the

AVA. These veterinarians have been responsible for the judicious use of antimicrobials since the 1980s (APV, 1989), and many of them have fostered and developed advanced technologies and management systems in pig production that have become standard practice on many farms. Any use of antimicrobials in the Australian pig industry can only occur under the guidance and direction of the farm veterinarian. The industry quality assurance program APIQ✓® underpins health management on most Australian pig farms. A formal Australian pork industry AMS plan is currently under development to consolidate and capitalise on AMS expertise, with the aim of supporting further improvements to practises that relate to AMS.

Australian Pork Limited (APL) has developed a number of supporting documents for optimal pig husbandry in Australia, including the Companion Handbook to the Model Code to support the Model Code of Practice for the Welfare of pigs (APL, 2013b). APL has also produced the 'Care of the Compromised Pig' manual (APL, 2013a) to assist producers in the early identification and treatment of pig health issues in addition to training through various registered training organisations and APL initiatives, which include training (at various levels) in identification, management and reporting of exotic and endemic diseases to minimise the risk of disease spread.

In addition, supervisors and staff working as stock-people on piggeries hold a Certificate III in Agriculture – Pig Production, or equivalent, which includes training for competency in maintaining the health and welfare of the pigs in their care.

Reduce

Many pig diseases are highly contagious and can be easily spread by a variety of vectors, and as most pig herds consist of pigs of varying ages, good biosecurity is critical. To this end, biosecurity is a stand-alone module in the industry's QA program, APIQ✓® (Module 4), and biosecurity performance indicators are also included in several other modules. New diseases or disease-causing agents of significant importance can be established in a herd through the introduction of new stock or semen to breed new stock, and biosecurity requirements also include management of these risks. The pork biosecurity manual is available at: <http://www.farmbiosecurity.com.au/industry/pigs>.

APL and its predecessor, the Pig Research and

Development Corporation, invested considerable resources to gain a greater understanding of the risks to the Australian pig herds' biosecurity (information available from annual reports at: <http://australianpork.com.au/library-resources/publications/annual-reports>). APL have also established a micro-site specifically dedicated to consolidating information regarding biosecurity and the maintenance of a high-health-status pig herd in Australia (<https://keepingourpigssafe.com.au>). The Australian pig industry's biosecurity endeavours have been greatly enhanced by the formation of the APL Biosecurity Strategic Review Panel, whose members are authorities in this field within, and outside of, the industry.

There are a small number of commercial breeder organisations in Australia that specialise in the development and distribution of commercial pork genetics within Australia. The Pork Cooperative Research Centre had an entire sub-program devoted to developing and implementing robust pork genetics, with a focus on breeding healthy, resilient and robust pig herds (further information can be found at: sub-program 2B; <http://porkcrc.com.au/research/program-2>). To further manage health and biosecurity, semen is often brought onto a farm to inseminate the sows for breeding, as opposed to bringing in new animals to a herd, which has a high risk of directly introducing disease.

Refine

The Australian pork industry has allocated resources through the High Integrity Australian Pork CRC and the APL to target specific areas focused on further refinement of antimicrobial use. Research outcomes have provided the platform for adoption of systems to improve management and, subsequently, the health and welfare of the pigs. The Pork CRC has been specifically investing in the development and refinement of diagnostic assays for Australian pigs (further information can be found at: <http://porkcrc.com.au/research/program-2>) and helping to implement these diagnostic assays in laboratories that support the pig industry and confirm disease treatment.

A recently completed APL project reviewed and standardised the efficacy of bacterial isolation and AMR methodologies in various laboratories that service the pig industry, to ensure that techniques and methodologies are consistent across the various labs so that comparisons can be made.

Many trading partners are either legislating, or have legislated, against the importation of pork products that have been produced with the use of growth promotants, however, their use has been phased out voluntarily in the

Australian pig industry and antimicrobials are only used to prevent and treat disease. Generally, the conditions that commonly require antimicrobials in pigs are enteric and respiratory diseases, which can lead to considerable pig morbidity and mortalities. In the Australian pig industry, there is a widespread reliance on drugs rated to be of low importance in the context of human health (Jordan et al., 2009), however renewed efforts are expected to reduce this dependence further.

Replace

Prior to the early 1980's, there were no vaccines available for some of the most common enteric and respiratory diseases in Australian pigs. Producers relied on antibiotic treatment and replacement fluids, and the efficacy of feed-back methods (a crude type of oral vaccination) was variable. Vaccine development for pigs was initiated in the early-mid 1980's and the impact on preventing disease was significant.

The Australian industry has access to commercial vaccines for many of the common pig diseases, including porcine circovirus associated disease, ileitis, parvovirus, leptospirosis, erysipelas, and the respiratory diseases enzootic pneumonia and pleuropneumonia (Appendix 3). Despite access to a vaccine, pleuropneumonia has proven difficult to control, even with the use of autogenous vaccines, and as such, there are ongoing investments to develop more effective pleuropneumonia vaccines and the identification of a vaccine candidate for swine dysentery. There are also ongoing investments in the development of more effective vaccines against bacterial and viral pathogens and the viability of alternative treatment options, such as the use of pre- and pro-biotics, bacteriophages, water/feed supplements, disinfectant fogging (to reduce bacterial loads in the environment) and organic acids that may not demonstrate antimicrobial action but improve gut health and function.

Different management systems have replaced traditional systems in an effort to better manage pig flow to reduce disease transmission between and within pig batches and housing cleanliness, and consequently, reduce antimicrobial use to treat bacterial diseases. These all-in/all-out systems of management can result in significant health improvements as they minimise transfer of pathogens from older to younger pigs, which assists in the maintenance of health status, however, these systems are not always economically viable and may not be adopted by all producers.

The voluntary industry quality assurance program, APIQ[✓]®, was developed and implemented in the 1990s and is the vehicle for many practises that safeguard the health and biosecurity of Australian pig herds. APIQ[✓]® certification allows pork producers to demonstrate their compliance with State/Territory and Commonwealth laws relating to food safety, animal welfare, biosecurity and traceability, and includes many elements of pig health. APIQ[✓]® has annual minor and major reviews every four years and export abattoirs only accept pigs that are APIQ[✓]® certified. APIQ[✓]® has transitioned to independent, or Third-Party, auditing. For an APIQ[✓]® certification to remain current, a producer's site/s must have an annual APIQ[✓]® Compliance Audit prior to the expiry of their certification, to have their APIQ[✓]® Certification approved for renewal. A significant aspect of compliance is an internal audit or review, which must be conducted approximately every six months and includes the review of on-farm manuals and records to ensure their systems remain current and are compliant.

Surveillance projects for the detection of AMR in bacteria from Australian pigs have been ad hoc and are summarised in the review by Abraham et al (Abraham et al., 2017). While only some AMR bacteria of concern have been identified, the results are favourable when compared internationally, and the industry is actively taking steps to address these issues. The elements of the pig industry's AMS program have already been voluntarily adopted by several producers and integrated into standard business operations. It is expected that the development of an industry wide AMS plan will assist the greater industry to undertake similar endeavours and identify further areas for research investments.



A group of brown chickens, likely a breed like the Brown Leghorn, are shown in a natural, straw-covered environment. The chickens have reddish-brown feathers and prominent red combs. They are positioned in the foreground and middle ground, looking in various directions. The background is filled with more chickens and straw, creating a sense of a busy farm or poultry house.

THE AUSTRALIAN POULTRY INDUSTRIES

THE AUSTRALIAN POULTRY INDUSTRIES

The Australian chicken meat, duck, turkey and egg industries are separate from each other, with their own industry representative organisations and RD&E arrangements.

The poultry meat industries are vertically integrated, and the farmers are predominately contractors to the processing companies who own the poultry. This relationship means that the processing companies are directly responsible for the farm inputs that are related to the poultry, including the feed, animal health, management advice, animal welfare standards and chicken harvesting. Flock health in the poultry meat industries is always managed by at least one registered veterinarian with expertise in poultry, who oversees and manages disease surveillance and prevention through biosecurity, vaccination and other integrated disease preventative strategies. The registered veterinarian is also responsible for disease diagnosis and treatment, including the administration of antimicrobials, for all contract, company and breeder flocks.

The scale of production in the Australian egg industry is incredibly varied and ranges from small backyard production to a few large producers and many small to medium-sized producers. In Australia, there are cage, barn and free-range egg production systems. Organic egg production is a niche segment contained within the free-range farms. Hens can lay at approximately 18 weeks of age and are in production until 80 weeks of age. Some larger egg producers also have a registered veterinarian with specialist expertise in poultry, who oversees and manages disease surveillance and prevention.

Very few countries have been able to meet the strict protocols for importing poultry products into Australia, and as a result, few poultry products are approved for importation into Australia. No live poultry or shell eggs that are intended for human consumption are imported into Australia, and any egg products that are imported are either preserved, cooked, pulped or in powder form.

Apart from chicken products imported from NZ, the only other poultry meat products currently imported into Australia are small volumes of processed chicken, duck or turkey meat products that have been fully retorted (i.e. cooked to high temperature in their container), such as in some canned foods, soups or animal feed. There are significant amounts of salted, cooked or preserved (“century”) duck eggs imported into Australia from Hong

Kong, China and the Philippines.

Poultry breeding stock for the chicken meat, egg, duck and turkey industries is imported as fertile hatching eggs that undergo strict quarantine protocols, which includes fumigation of the eggs and testing for various pathogens in the overseas donor flocks. The fertile eggs are hatched out in quarantine stations in Australia and the chicks hatched from them are subject to rigorous testing for key pathogens before being released to highly biosecure breeder farms.

The Australian chicken meat industry is dominated by six companies that supply the bulk (approximately 95%) of domestically produced chicken meat. Meat chickens reach processing age at between 5 – 7 weeks of age, in either barn or free-range production systems, and all chickens are grown on the floor of large barns that are either climate-controlled or naturally ventilated. The chicken meat production system is ‘all-in/all-out’, which means there is a period between flocks when the sheds are emptied, cleaned and sanitised. This breaks the pathogen cycle and minimises the proliferation of bacteria and transfer to subsequent flocks.

The two largest meat chicken processors also produce the bulk of Australian turkey meat, alongside several smaller turkey companies that breed, grow and process their own flocks. Turkeys reach processing age from eight weeks of age.

The Australian duck meat industry is dominated by two large companies and two smaller enterprises, which supply the majority of domestically produced duck meat. Ducks are raised to approximately 7 weeks of age on the floor of large barns that are usually naturally ventilated with some tunnel-ventilated.



Responsibility

Antimicrobial stewards for the poultry industries are the veterinarians employed by, or working with, the poultry companies that operate in Australia. Australian registered veterinarians that have specialist expertise in poultry and are engaged in the management of the commercial poultry industries are members of the Australasian Veterinary Poultry Association (AVPA) and/or the Commercial Poultry Veterinarians, the latter being an AVA special-interest group. The Australian poultry veterinarians have a long, documented history of embracing the principles for the appropriate use of antimicrobials. Their most recent codes of practice, 'The code of practice for the use of antibiotics in the poultry industry' (AVPA, 2001), was last reviewed in 2001, but the original version was produced in 1987 (APIA, 1987). The next review of this document will be undertaken with consideration given to reflect any potential changes to AVA documents, which are also planned for review, along with the development of the AVA prescribing guidelines for poultry.

The poultry industries and the AVA jointly developed a "Code of Practice for the Use of Prescription Animal Remedies (Schedule 4 Substances) in the Poultry Industry" (further information can be found at: <http://www.ava.com.au/policy/26-code-practice-use-prescription-animal-remedies-schedule-4-substances-poultry-industry>), which was ratified by the AVA in 2005.

In 2018, the 'Model Code of Practice for the Welfare of Animals: Domestic Poultry' (PISC, 2002), is being revised and rewritten into Standards and Guidelines (expected to be completed by 2019), which includes requirements for adequate husbandry and disease prevention strategies to reduce the incidence of poultry diseases in Australian poultry flocks.

Poultry breeders

There is clear recognition that the use of antimicrobials in flocks and hatcheries at the top of the 'breeding pyramid' may affect the characteristics of the microbiological populations of subsequent, much larger, generations of poultry.

The operations involved in supplying poultry genetics to the commercial Australian market are either owned by, or directly associated with, international breeding companies. The primary poultry breeding operations in Australia employ veterinarians that are directly involved in the overall management of the breeding stock and who are responsible for prescribing and overseeing the administration of all antimicrobials used

in poultry breeding operations. Veterinarians are given independence in decision-making related to antibiotic use and are supported by local and international company management.

Poultry meat industries (chicken, duck and turkey)

Due to similarities in the vertically integrated structure of the chicken meat, duck meat and turkey meat operations, there are a small number of skilled veterinarians responsible for the flock health programs and stewardship in the poultry meat industries. This has allowed, and continues to facilitate, prompt improvements made to poultry management when new evidence is supplied to support change in practices. As part of the contractual arrangements, poultry meat farmers are required to adhere to company specified practices (these are commercial-in-confidence) and there are often financial penalties to farmers whose flocks experience an increase in mortality or morbidity. Flock supervision and monitoring must adhere to company policies, procedures and industry manuals, with implementation and progress monitored by company representatives, including the veterinarians, during frequent farm visits. In-house training is provided for farmers and company service personnel and livestock managers on early disease identification and optimising husbandry and biosecurity for each farm.

In 2017, the chicken meat industry formalised an AMS plan that was implemented in the companies that produce approximately 95% of the chicken produced in Australia. This plan was tailored to operations within each of the companies and will be independently verified in 2019, with the aim of identifying further areas for improvement in AMS.

The egg industry

There are a few large producers in Australia that supply the bulk of fresh egg products consumed domestically and as a result, there a small number of skilled poultry veterinarians responsible for the health and stewardship of antimicrobials in layer hens.

However, for the niche and small-to-medium sized producers, access to a poultry veterinarian may not always be possible or affordable, which often means they access companion animal veterinarians or other medical support to manage the health of their flocks, even though only registered veterinarians can prescribe antibiotics.

Australian Eggs has developed training programs for farm staff, which include consideration for health, welfare and biosecurity; the essential parts of AMS. An induction training program ('Eggstart') was established in 2016 and a Certificate III in Poultry Production was established in 2011, both of which are coordinated through a third-party training organisation. Australian Eggs Ltd is also in the process of formalising an AMS plan for the industry, which is expected to be available for implementation in 2019.

Reduce

Prevention of disease in poultry flocks is paramount. Since the 1980s, there has been ongoing work to better understand poultry diseases, define biosecurity controls, improve precision of nutritional management and develop and implement extensive flock vaccination programs (Appendix 3). This has resulted in a significant reduction in the incidence of respiratory and gastrointestinal diseases that could otherwise require treatment or prevention with antimicrobials. As a consequence, there has been a reduction in the overall use of many antimicrobials in poultry production. The recent growth in free-range and non-barn/cage farming practises means that there is an increase in poultry with greater exposure to environments that may contain pathogens, which has resulted in the re-emergence of some diseases that sometimes require treatment with antimicrobials, when other preventative or treatment options fail or aren't available.

The National Farm Biosecurity Manual for Poultry Production outlines the minimum recommendations for biosecurity that capture risks relevant to all poultry operations (further information can be found at: <http://www.farmbiosecurity.com.au/livestock/>), and the chicken meat, egg and duck industries have each developed industry specific biosecurity manuals that are currently under review.

Disease prevention, through biosecurity, optimum nutrition, vaccination and improved husbandry procedures have been, and continue to be, key to ensuring production of healthy poultry without being excessively cost prohibitive.

Primary poultry breeders

Whenever possible and viable, methods of reducing the use of antimicrobials have always been assessed and implemented by poultry breeders. There is an emphasis on stringent, audited biosecurity throughout the poultry primary breeding industry internationally, and particularly in Australia where all new poultry genetics is imported. This aims to prevent the introduction of exotic diseases

and endemic agents that may cause disease within valuable genetic flocks, which can significantly impact on the availability of commercial poultry for production. Access to primary breeding operations is strictly controlled and all material inputs undergo risk assessment before coming into contact with flocks. Through biosecurity and flock monitoring, the primary Australian breeding industries have become free of agents that had previously been controlled with antimicrobials, such as *Mycoplasma* species, which is considered a significant achievement.

Hygiene on farms and in hatcheries must be of a standard that can reduce the contamination and challenges encountered by newly hatched chicks. For example, hatching egg hygiene on-farm must include clean nest boxes, dry, friable litter, clean egg belts, hand washing, and clean and disinfected storage facilities to ensure the highest hygiene standards in the hatchery. Fumigation of eggs, cleaning and disinfection of hatchery equipment and personal hygiene of staff that handle newly hatched chicks are all standard in primary breeding hatcheries. In 2017/18, two of the largest primary breeding companies in Australia invested heavily in new hatchery buildings and technology, with the aim of improving hygiene and reducing early chick mortality.

Continual improvements in the genetics of poultry breeds are controlled and undertaken by international nucleus breeding companies, to produce an immunocompetent, robust hybrid bird at the commercial production level. This is coupled with refined management practises to significantly reduce the incidence of disease and, therefore, the need for antimicrobials over time.

Poultry meat industries (chicken, duck and turkey)

Poultry meat farmers are required to adhere to the relevant National Biosecurity Manual as part of their contractual obligations with the processing companies and various auditing requirements.

Commercial operations have always continually assessed operations to identify areas for improvement in biosecurity and disease prevention, in order to ensure healthy and productive poultry. This has reduced the need for treating, or routine prevention of, illness in the poultry meat industries with the use of antimicrobials. This is particularly important as the structure of the industries changes to consolidate operations into fewer, larger farms.

The evolution of improved duck drinking systems (nipple drinkers) and the reduced use of troughs and surface water has had a profound impact on the incidence of bacterial disease in growing ducks, with similar impacts on fertile egg hygiene and embryonic survival. High standards of endemic disease control have reduced the number of significant disease incursions since 2012.

The egg industry

Improvements in husbandry and continual upgrades in housing have reduced the need for antimicrobials in egg production. However, the recent growth in free-range and non-cage egg farming has resulted in the re-emergence of some diseases, such as coccidiosis, fowl cholera, erysipelas and spotty liver disease (SLD), which has been identified as caused by a new aetiological agent, *Campylobacter hepaticus* (Van et al., 2016). These diseases require treatment with antimicrobials where relevant, and when other preventative or treatment options are not available, or appropriate.

Refine

For most of the poultry meat produced in Australia, routine diagnostic testing is performed to inform veterinary treatment of flocks that are demonstrating clinical signs of bacterial disease. More than 25 molecular and/or diagnostic assays are available at private, in-house and public veterinary diagnostic laboratories to support the diagnosis of avian diseases present in Australia (an example list of diagnostics can be found at: https://fvas.unimelb.edu.au/__data/assets/pdf_file/0010/2860804/APCAH-Customer-Commercial-Price-List-PCR-and-Serology-FEB-2018-v.2.pdf). There is extensive collaboration across the poultry industries for on-going investments in research to provide rapid, effective typing for avian diseases.

As part of the commitment to judicious antimicrobial use, the Australian poultry industries reviewed the efficacy of AMR methodologies in the 1980s to ensure that AMR detection and surveillance approaches are consistent (Whithear et al., 1986). Since then, there has been greater alignment of techniques used by industry laboratories for detection and reporting of AMR, although improvements can still be made. As such, current RD&E investments aim to further align the bacterial isolation and characterisation methodologies across public and private laboratories that support the poultry industries.

In 2018/19, a research project funded by the Australian Government Department of Agriculture and Water Resources, aims to align bacterial isolation and AMR

methodologies in laboratories that service the poultry industry. This will help determine whether current industry resources can be utilised as a framework for ongoing industry-level AMR surveillance needs.

Primary poultry breeders

The use of antimicrobials in the Australian primary poultry breeding industry is tightly controlled and can only be administered after investigation and diagnosis by a registered veterinarian. When possible, medication will only be administered after a culture and sensitivity assay is performed to ensure the antimicrobial used will be effective. Records of all treatments are completed.

Poultry meat industries (chicken, duck and turkey)

The Australian Chicken Meat Federation (ACMF) established a policy in 2007 that antibiotics should not be used for growth promotion purposes, and has been actively working with the product registrants since then to have growth promotion claims for chickens removed from labels (ACMF, 2018b). There are currently two products that remain registered for use in poultry that have growth promotion claims, however, neither of them are used in human medicine. One of them is not available for sale in Australia, and the other is occasionally used to treat enteritis when other preventative and treatment measures have failed to control disease.

The chicken meat and turkey meat industries generally use in-feed medication for the prevention of common diseases, such as coccidiosis and necrotic enteritis, which are highly likely to be present in a flock. Water medication is generally used for treatment of diseases post-diagnosis by a veterinarian. Due to the short production cycle, there are regular opportunities to trial refinement of preventative therapies, whether using antimicrobials or other products.

Disease prevention and refinement of antimicrobial usage are the predominant drivers to producing healthy turkeys without being excessively cost prohibitive. Focus on hygiene, management and biosecurity are key underlying measures to prevent disease in turkeys, and the industry continually refines the use of the few tools available to support health management in turkey flocks.

The availability of autogenous vaccines for *Riemerella* and *Pasteurella*, plus the adoption of dry production systems with good shed litter management, has enabled antimicrobial use to be almost eliminated in duck production in Australia. Treatments are generally only used in response to a disease outbreak and in these cases, the limited therapeutic treatments for ducks are conventionally provided by water medication. For most duck meat produced, routine diagnostic testing is performed to inform veterinary treatment of flocks displaying clinical signs of disease. Antimicrobial susceptibility testing is undertaken where available.

The poultry meat industries endeavour to continuously assess gaps in disease prevention and determine areas where improvements in biosecurity can be made to reduce the introduction and spread of pathogens. These industries have made, and continue to make, investments into improving biosecurity with the aim of reducing the introduction of pathogens and therefore the need for treatment or preventive uses of antimicrobials to manage disease in meat poultry flocks.

The egg industry

There are different antimicrobials that can be used for prevention or treatment of disease depending on whether the hens are laying or are in the weeks prior to coming into lay (termed 'pullets'). The reason for this is to prevent potential antimicrobial residues in the eggs, as only antimicrobials with a nil withholding period can be used in layers during lay.

Effective biosecurity, nutrition, vaccination and husbandry are crucial to ensuring a high health status for hens to reduce the number of preventative treatments required. The egg industry actively supports ongoing projects in collaboration with the APVMA and other regulatory authorities, to allow the introduction of medications that are widely used internationally but not currently registered for use in the Australian layer industry. This is to ensure the health and welfare of hens and may help refine, or remove, the use of antimicrobials that are of importance to human medicine.

Replace

The Australian poultry industries have a long history of collaboratively identifying, developing and implementing alternatives to antimicrobials. There is increasing use of alternative therapeutic options, such as probiotics, prebiotics, organic acids and other feed supplements that aim to improve gut health by modifying the intestinal microbiota. However, the focus for disease prevention has

historically been, and continues to be, high standards of biosecurity coupled with the use of vaccines.

Vaccines available in Australia for the prevention of poultry diseases are outlined in Appendix 3, along with *Riemerella* and *Pasteurella* vaccines in ducks. Autogenous vaccines are also used. Vaccine use is dependent on the risk of infection in the region where the poultry are raised, and in some cases, an industry may vaccinate every chick hatched nationally (e.g. Marek's Disease and Infectious Bronchitis in the egg industry). Vaccines are applied either directly to commercial poultry or to breeder stock, to build adequate early immunity and to break the cycle of infection. Many vaccines are used in the primary breeding stock due to the high nationally significant economic repercussions of disease in these flocks.

Prior to the development and implementation of many of the currently available poultry vaccines, chronic respiratory disease complexes were the greatest source of economic loss due to disease in poultry, along with the secondary bacterial infections that required treatment with antimicrobials (Delaplane and Stuart, 1943; Bagust, 1989; Mifflin and Blackall, 1998).

In response to this challenge, the Australian poultry industry funded the development of vaccines to prevent mycoplasma-associated diseases (Whithear et al., 1990), which has meant macrolide and pleuromutilin antibiotics are rarely indicated or required in Australian poultry industries. These vaccines are now used internationally. The Australian poultry industries have recently funded the development and licencing of a vaccine candidate for necrotic enteritis (caused by *Clostridium perfringens* infection), which looks promising and, if it results in a commercially viable product, implementation would significantly reduce the need for some preventative antimicrobial use in Australia and internationally.

The poultry RD&E and industry representative organisations work with the vaccine supply companies through various industry and State government groups, to ensure there are viable and sustainable options available for preventing disease in poultry. There are also additional candidate vaccines currently going through the commercialisation and registration process, and continued investment from the R&D funding organisations and veterinary vaccine companies into improving the efficacy of current vaccines.

Poultry meat industries (chicken, duck and turkey)

Improved vaccination strategies have reduced antimicrobial use over time, as they help to protect more poultry more effectively, through the prevention of immunosuppressive diseases (such as Marek's disease, chicken anaemia virus and chicken infectious bursal disease) and the accompanying secondary bacterial infections or vertically transmitted diseases (e.g. Mycoplasmosis). Improvements in biosecurity and hygiene in these industries have not only reduced the use of antimicrobials, but in some instances, have also reduced on the need for vaccination.

Vaccination (particularly autogenous vaccines for Riemerella and Pasteurella) has been very successful in controlling almost all the important bacterial diseases that cause disease in Australian intensive duck systems, which is considered to have reduced the use of antimicrobials (particularly amoxicillin and oxytetracycline) to almost zero.

Implementation of tighter biosecurity controls, precise nutritional management and the development and use of Pasteurella and Erysipelas vaccines has resulted in a reduction in the need to use antimicrobials in the Australian turkey industry. Vaccination is widely applied where available, and new vaccines for meat chickens are often trialled for efficacy in turkeys, although this is not always successful. While there is significant R&D undertaken in turkey production (including vaccine development) in the USA and other countries, many of those advances cannot be adopted in Australia, usually due to import biosecurity restrictions. The Australian poultry industry also lacks the relevant infrastructure to undertake significant R&D for alternatives to antimicrobials for turkeys, and resources are limited as there is no R&D levy on turkeys (which means there is no formal R&D program). Despite this, progress is being made to develop an Australian-turkey specific *Mycoplasma gallisepticum* vaccine, and a haemorrhagic enteritis vaccine is expected to be commercially available by 2020 (Clements, 2015). In the meantime, there is continuous assessment of other alternative treatment options, such as the use of pre- and pro-biotics and feed supplements that do not have direct antimicrobial action, but which may improve gut health and function and be effective replacements for antimicrobial use in turkeys.

The egg industry

The increase and expanding number of hens farmed under extensive conditions, such as free-range, has resulted in the re-emergence of diseases (such as parasitic infections and bacterial diseases caused by

Pasteurella). These diseases were better prevented in more intensive systems where the faecal oral cycle was limited (e.g. cage), however, the limited availability of antimicrobials to use in hens, particularly during lay (due to the concern of antimicrobial residues in eggs), means there is a greater reliance on disease prevention strategies.

Since 2016-17, the egg industry has heavily invested in the development of vaccine candidates to minimise the impact of Spotty Liver, which is a new emerging disease (Van et al., 2017).

Review

Given the long history of interrelationships between veterinarians in the poultry industry through the AVPA, there is regular opportunity to review industry level efforts and identify opportunities for improvement. This has resulted in harmonisation across the poultry sectors in relation to the adoption of AMS principles in practise.

Surveillance projects for the detection of AMR in bacteria from Australian poultry have been ad hoc, with most of the public information available for the chicken meat and egg industries, as they are the two predominant poultry industries in Australia. Routine antimicrobial susceptibility testing is considered commercially sensitive and, subsequently, is not reported outside the companies. Prior to the AMR surveillance studies funded by the Australian Government for the egg and chicken meat industries in 2016/17, the chicken meat industry funded RD&E to assess the levels of resistance in bacteria isolated from chickens in Australia in 2000/01 (Barton and Wilkins, 2001). Results to date indicate that the presence of AMR bacteria in Australian poultry is low in comparison to other countries.

Primary poultry breeders

The small and very valuable nature of the primary poultry breeding industry in Australia means there is tight managerial and veterinary oversight of all flocks. Health issues must be, and are, identified and addressed quickly. Efficacy of treatments is continually assessed and subsequently modified by a process of continual improvement. Despite no formal QA program for the poultry breeder industry, audits are conducted by the Australian Government of Agriculture and Water Resources to approve sites for export, and some are ISO and HACCP accredited. Due to the high value of primary breeding stocks, extensive records are kept of husbandry practises, including medications that may be administered, throughout the life of the flocks.

Poultry meat industries (chicken, duck and turkey)

There are no national QA Programs for the chicken meat, duck meat or turkey meat industries. This means that each company has different approaches and requirements based on their own situation and their customers' needs, including customer (e.g. Coles, Woolworths, McDonald's etc) programs and third-party QA accreditation schemes (e.g. FREPA [Free Range Egg and Poultry Australia]; RSPCA). All these programs have strict requirements for keeping accurate and detailed production records, including details of medications provided. Records and industry standard practices are subject to separate audits for all required accreditations and programs on a regular basis.

As there is no formal industry-led method for auditing and verification, the industry-led AMS program that was adopted voluntarily and integrated into standard business operations for the chicken meat industry in 2016/17 requires independent verification, which is planned to be undertaken in 2018/19. It is expected that several recommendations for future research to support industry AMS will be identified as part of this process.

The egg industry

The Australian egg industry has a voluntary industry QA program, Egg Standards of Australia (ESA), which was implemented in 2016 (superseding the Egg Corp Assured, which commenced in 2009) and provides a framework to demonstrate compliance against a set of standards for quality egg production in Australia. ESA has been developed to provide a compliance framework for egg farmers to meet the needs of regulators, retailers, farmers and egg buyers in areas including hen welfare, egg quality, biosecurity, food safety, work health and safety and environmental management. Audits are conducted at least once per year. The ESA program itself is regularly reviewed to ensure they are up-to-date with current knowledge on disease prevention, treatment and husbandry, and relevant supporting materials, including those for AMR/AMS, are available to support producers.

Under ESA, medications used must be recorded and the principles of AMS are currently being formally incorporated to assist vets and farmers in recording all details regarding the use of antimicrobials, including a record of script and use directions from a veterinarian. Given the wide variation in the size of egg businesses, not all are accredited under ESA, however, the companies that produce the bulk of eggs and egg products sold commercially are ESA accredited.





THE
AUSTRALIAN
RED MEAT
INDUSTRIES

The Australian Red Meat Industries

The Australian sheep meat and grain-fed and pasture-fed cattle industries are separate from each other with their own industry representative organisations, although all R&D for these industries is coordinated through Meat & Livestock Australia (MLA) as their RDC. The Australian cattle and sheep meat industries focus on meeting consumer expectations, conducting research to ensure a sustainable future, and encompass scientific approaches to production and disease management. The industries have a broad geographic distribution across Australia, with a wide range of production sizes and thousands of producers.

In 2017, Australia was ranked as one of the world's most efficient producers of cattle and one of the largest exporters of beef. As of 2017, Australia is also the world's largest exporter of sheepmeat and is the world's second largest producer of lamb and mutton (MLA, 2017).

Most cattle and sheep are the product of extensive grazing systems, where they spend most of their lives on pasture eating grass with additional supplementary feed when necessary. Grain-fed cattle spend 70 – 100+ days in a feedlot to improve body condition and enhance the quality of beef meat with a controlled environment. Similarly, a small number of sheep may spend a short time in a feedlot to improve and standardise their meat quality or help when seasonal conditions dictate the need for this system. Some cattle may be kept in a feedlot for an extended period to produce meat of exceptional quality, such as highly marbled beef. Cattle and sheep reach slaughter age anywhere between 12 – 30 months, and potentially much earlier or much later as slaughter age is often based on maturity/production definitions rather than chronological age. Importation of beef (chilled or frozen) and beef products is permitted from Japan, the Netherlands, New Zealand, the United States and Vanuatu, and strict import protocols mean no live cattle are imported into Australia. While theoretically it is possible to import live sheep into Australia from New Zealand, the import requirements make it commercially difficult and likely unviable. There have been no live sheep imports from New Zealand since 2016.

Veterinarians are often hired to aid with animal health management and are required for prescribing medications (including antimicrobials), and feedlot veterinarians and other staff inspect cattle every day to identify early signs of disease. Few cattle receive antibiotics at any time during their lives in Australian extensive agriculture. Anecdotally, it is considered that there would be, “far less

than one course of antimicrobials used per animal per year” (MLA, 2014). Due to the low usage of antimicrobials, the presence of antibiotic resistant bacteria is also low (Barlow et al., 2015, 2017).

Responsibility

The antimicrobial stewards in the red meat industries are the prescribing veterinarians and the farm or feedlot team who are responsible for good animal care practises, including infection control and prevention. A significant proportion of veterinarians with expertise in livestock support the red meat industries and many of them are members of special interest groups of the AVA (Australian Cattle Veterinarians [ACV] and the Sheep, Camelid and Goat Veterinarians [SCGV]), which provide opportunities for professional networking and discussion about current practise, as they are all supported by various AVA policies and guidelines. In 2014, MLA produced a factsheet that outlined the AMR status and AMS initiatives in the cattle industries in Australia to that date (further information can be found at: <https://www.mla.com.au/globalassets/mla-corporate/research-and-development/program-areas/food-safety/pdfs/antimicrobials-and-the-cattle-industry-fact-sheet.pdf>).

The Australian Lot Feeders Association (ALFA) launched the comprehensive ‘Antimicrobial stewardship guidelines for the Australian cattle feedlot industry’ in early 2018 (<https://www.mla.com.au/research-and-development/animal-health-welfare-and-biosecurity/diseases/>), which outlines the AMS plan for the industry and is now in the implementation phase (MLA, 2018a). The report also supplies supporting tools, such as antimicrobial usage calculations, guidance on how to produce an AMS plan and recommendations for lot feed producers (MLA, 2018a). These guidelines provide a continuous improvement framework that will help feed-lot producers understand and ensure their appropriate use of antimicrobials is aligned with the National AMR Strategy (Australian Government, 2015). An extensive array of tools and training opportunities for cattle are also provided by MLA (further information can be found at: <https://www.mla.com.au/extension-training-and-tools/>).

The model code of practice for the welfare of cattle in Australia was converted to Standards and Guidelines in 2016, which are in the process of being implemented into state welfare legislation. The report includes requirements

for the person in charge to have knowledge of, and manage, the health and welfare of the cattle in their care (further information can be found at: <http://www.animalwelfarestandards.net.au/cattle/>).

Reduce

Commercial industries have always strived for continuous improvements in the understanding and management of biosecurity, husbandry, hygiene, cattle and sheep physiology and quality nutrition, as these issues are central to preventing disease, and the related productivity issues. The Livestock Production Assurance program has 215,000 participants and part of this program requires producers to have a Farm Biosecurity Plan and integrate best-practice biosecurity in their on-farm management. This program was relaunched with enhanced requirements in 2017 and further information can be found at: <https://www.mla.com.au/meat-safety-and-traceability/red-meat-integrity-system/about-the-livestock-production-assurance-program/>.

The National Farm Biosecurity Manual for Grazing Livestock Production and a significant list of additional support tools and resources for biosecurity have been produced in consultation with industry and government agencies for the cattle (<http://www.farmbiosecurity.com.au/industry/beef-cattle/>) and sheep meat (<http://www.farmbiosecurity.com.au/industry/sheep/>) industries. There is also a Farm Biosecurity Manual specifically for cattle held in feedlots (<http://www.farmbiosecurity.com.au/industry/lot-feeding/>) and biosecurity requirements are captured as part of the feedlot industry's accreditation program, the National Feedlot Accreditation Scheme (<https://www.ausmeat.com.au/services/list/livestock/nfas/nfas-information/>) (https://www.ausmeat.com.au/WebDocuments/NFAS_-_Rules_and_Standards_of_Accreditation.pdf). This program requires feed-lot producers to undertake a risk assessment and implement a Biosecurity Management Plan. The red meat industries also partner with Animal Health Australia to establish and coordinate various biosecurity initiatives to provide red meat producers with the necessary tools and useful information to manage disease (endemic and exotic), pest and weed events on their farms.

MLA invests up to \$5 million a year in livestock genetics R&D under a standalone R&D program, to facilitate increased genetic gain and health robustness for cattle and sheepmeat industries. A number of resources and research reports have been developed to assist producers in genetic selection and breeding in cattle and sheep (further information can be found at: <https://www.mla.com.au/research-and-development/Genetics-and-breeding/cattle/genetics/>;

<http://mbfp.mla.com.au/Cattle-genetics/>; <https://www.mla.com.au/research-and-development/Genetics-and-breeding/sheep/>; <http://www.sheepgenetics.org.au/Home>).

Cattle in feedlots are considered more susceptible to ailments caused by seasonal influences that may require antimicrobial treatment. Several research studies are underway to support previous work related to reducing the use of antimicrobials in feedlot cattle and other red meat industries (further information can be found at: <https://www.mla.com.au/research-and-development/animal-health-welfare-and-biosecurity/>). These projects range from investigating better ways of preparing cattle to enter a feedlot (for example, becoming used to being in larger social groups), determining the best vaccination regimen for cattle prior to feedlot entry, and faster diagnosis of infection in cattle while in a feedlot.

The grazing cattle and sheepmeat industries have low rates of infectious disease because of inherently low stocking rates and the use of preventive measures, including vaccination, stock handling practises, insect control, biosecurity, herd management and infection control (Barnes, 2015; MLA, 2014), which has resulted in significant reductions in the use of antimicrobials in these industries over time and is reflected in the low frequency of AMR.

Under its 2015-20 Strategic Plan, the sheepmeat industry is working to reduce the cost of endemic diseases, including internal and external parasites, by \$3 million



in 2020 and \$69 million by 2030. This cost saving to the industry is expected to be predominately due to a reduction in illness, reduction in the need for antimicrobial treatments and the loss of productivity from sick animals.

Refine

The use of antimicrobials in red meat industries is guided by veterinary prescribing guidelines such as those issued and reviewed by the AVA (AVA, 2017). An extensive suite of diagnostics has been developed and are regularly used by veterinarians to diagnose disease and identify appropriate treatment (further information can be found at: <https://www.dpi.nsw.gov.au/about-us/services/laboratory-services/veterinary/veterinary-test-list-by-species/cattle>). MLA invests millions of dollars each year in improvements to diagnostics and disease control (further information can be found at: www.mla.com.au/research-and-development/animal-health-welfare-and-biosecurity), which inform the industry's best practice manuals that underpin the various accreditation programs in place, such as the National Feedlot Accreditation Scheme and Livestock Production Assurance programs (further information can be found at: <https://www.ausmeat.com.au/services/list/livestock/nfas/nfas-information/>; https://www.ausmeat.com.au/WebDocuments/NFAS_-_Rules_and_Standards_of_Accreditation.pdf; <https://www.mla.com.au/meat-safety-and-traceability/red-meat-integrity-system/about-the-livestock-production-assurance-program>).

Replace

Australia is fortunate to be free of all the major epidemic diseases in red meat livestock and is relatively free of other serious parasites and diseases (AHA, 2018). Of the endemic diseases that do affect these industries, parasitic diseases have the largest financial impact on farm productivity.

Vaccines that are available in Australia to help prevent common endemic diseases in red meat livestock are listed in Appendix 3. These include vaccines to help prevent Bovine respiratory disease, Clostridial diseases (including botulism), Cheesy gland, John's disease, Leptospirosis, Pestivirus, Pinkeye, Three-day sickness and Vibriosis. These diseases can have a significant impact on the health, welfare and productivity of livestock. Vaccines are used at different times in the production cycle depending on the disease, the vaccine and the risk of infection in the region where the cattle or sheep are being produced.

The replacement of antimicrobials of importance to human medicine is considered whenever evidence becomes available that supports the efficacy and safety

of an alternative. Adoption is carefully assessed to avoid unintended consequences of harm to the animals, or subsequent increased need for the use of antimicrobials of importance to human medicine if the vaccination fails.

There is significant effort underway to identify and assess antimicrobial alternatives through various MLA-funded programs. Currently, there are few alternatives to antimicrobials for treating unwell red meat livestock, and the best options for reducing antibiotic use lie in the prevention of infection through the use of vaccines, good biosecurity and hygienic husbandry. Extensive resources to support the health and welfare of red-meat livestock are available from MLA, as well as the numerous other organisations that support the red-meat industries (further information can be found at: <http://mbfp.mla.com.au/Herd-health-and-welfare>).

The use of antimicrobials in beef, particularly pasture-fed, and sheepmeat has always been low, and often for practical reasons, a farmer will avoid calling upon a veterinarian for consultation and treatment unless necessary. As a result, preventative measures will always be preferable to treating disease.

Review

The Livestock Production Assurance (LPA) program was introduced in 2004 and is the Australian red meat industry's on-farm program covering food safety, animal welfare and biosecurity. It is part of the integrity system used by the red meat industry to meet the stringent requirements for domestic and export markets (further information can be found at: <https://www.mla.com.au/meat-safety-and-traceability/red-meat-integrity-system/about-the-livestock-production-assurance-program/>).

More than 200,000 producers are part of the LPA, which covers requirements for safe and responsible animal treatments, stock foods, fodder crops, grain and pasture treatments, biosecurity and animal welfare, and includes numerous learning modules for accredited producers. It is a requirement of LPA that stringent records are kept of how the meat was produced, including detailed records of medications administered. This information is used for analysis of use and efficiency to guide future decisions on treatment protocols, antimicrobial use and to allow quality of use to be assessed. However, given the structure of the industry, this information is not easily captured at a national level.

Due to the significant reliance on export markets for the red meat produced in Australia, the industries maintain surveillance of developments in AMR and stewardship initiatives and conduct research to measure its

performance against recommendations. MLA has funded several surveillance projects in the Australian red-meat industry (Barlow et al., 2015, 2017), with another planned for 2018/19, and results to date demonstrate very low levels of AMR in bacteria from cattle in Australia.

Grain-fed beef industry

The Australian Lot Feeders Association implemented the National Feedlot Accreditation Scheme in 1994, which was the first QA program implemented in Australian agriculture (further information can be found at: <http://www.feedlots.com.au/industry/nfas>). This QA program was comprehensively reviewed in 2015 (Cudmore JW, 2015). It requires every accredited feedlot to be independently audited on an annual basis, to ensure they comply with animal health, welfare, environmental and food safety legislation. As of 2018, 400 feedlot producers are accredited through NFAS, which is a voluntary industry self-regulatory, QA scheme, although accreditation is mandatory for feedlots that produce for export markets. To be accredited under the NFAS, feedlots must have procedures that meet the requirements of industry standards and maintain records to demonstrate that these procedures have been adhered to for all cattle prepared at the feedlot (MLA, 2018b). The procedures, records and feedlot facilities must then undergo a third-party audit and feedlots must have a specific number of QA officers at the feedlot (number on site is related to the size of the feedlot; further information can be found at: <https://www.ausmeat.com.au/services/list/livestock/nfas/nfas-information/>).

In what is believed to be a first for feedlot industries around the world, the Australian industry implemented the Antimicrobial Stewardship Guidelines (MLA, 2018a) in 2018, to help guide judicious use of antimicrobials without compromising animal health. The AMS program is planned for regular review to allow for the adoption of processes for continual improvement. This will be used to evaluate compliance with the initiatives and to ensure that if antimicrobials of importance to human health are required to be used, their use reflects contemporary best practice. A review of the status of animal health, wellbeing and antimicrobial use are captured as part of this program. During the review phase of the AMS program, areas in need of improvement are identified and a plan was

devised to drive improvement on individual operations. The outcomes are monitored and documented and include measurements of quantity and quality for each use of an antimicrobial to capture appropriateness of antimicrobial use (i.e. has the treatment protocol been followed).

Sheepmeat industry

In extensive agriculture, few sheep ever receive antibiotics at any time during their lives. Sheep Producers Australia has recently initiated two projects that further underpin the sheepmeat industry's commitment to food safety and responsible use and management of antibiotics. The Agvet Chemicals in the Sheepmeat Industry project will determine potential risks of residues and contamination, and the management of risks associated with contamination of sheepmeat products that Agvet chemicals may impose. The Sheep Feedlots – Identification and Management of Risk project will determine potential risks of residues and contamination, and the management of risks associated with contamination of sheepmeat products that the sheep feedlot sector may impose.



iStock



THE
AUSTRALIAN
DAIRY
INDUSTRY

THE AUSTRALIAN DAIRY INDUSTRY

The national dairy cow herd in Australia is 1.56 million cows, with an annual milk production per cow of approximately 6070 litres and a total annual milk production of 9,289 million litres. 'Cheese' (36% milk utilisation; 377,727 tonnes produced annually), 'drinking milk' (27% milk utilisation; 1,423 million litres annually), 'milk powders' (23% milk utilisation for skim milk powder or butter; 273,425 tonnes annually) and 'butter' (92,698 tonnes annually) are the main commodities produced. Approximately 36% of Australian milk production is exported (<https://www.dairyaustralia.com.au/Industry/Dairy-Situation-and-Outlook/Situation-and-Outlook>; Dairy Australia, 2018).

History of antimicrobial stewardship in the Australian dairy industry

Antimicrobial agents play an important role in ensuring the health in the Australian dairy herd. They are used to prevent and control bacterial diseases in populations in order to protect the health, welfare and productivity of Australian herds. Although the dairy industry is committed to ensuring access to effective animal health treatments, it is very much aware of the consequences of their use and the potential for the development of resistance, severely compromising effective use in humans and animals.

The International Dairy Federation (IDF) have produced a factsheet for judicious use of antimicrobials for the global dairy industry. Dairy Australia was a major contributor to the development of this guide (available here: https://www.fil-idf.org/wp-content/uploads/2017/05/Factsheet-003_2017-Guidance-on-Antimicrobial-Resistance-from-the-Dairy-Sector.pdf).

Although antimicrobial usage and consumption appears to be relatively low in Australian agriculture, compared to the rest of the World the Australian dairy industry has implemented several strategies that provide guidance on, and improve implementation of, antimicrobial stewardship principles particular relating to improvements in farm biosecurity, hygiene practises and animal husbandry (for examples see: <http://www.sustainabledairyoz.com.au/>; <http://drycowconsult.com.au/>; <https://www.dairyaustralia.com.au/healthycalves>).

Australian dairy farmers are privileged to have an extensive support network who help maintain healthy cows and milk, which reduces the need for antimicrobial treatments. This network includes factory field officers,

milking machine technicians, veterinarians, dairy extension specialists and farm consultants who work with dairy farmers to improve the health of dairy animals.

Many programs have been developed by the Australian dairy industry to reduce the requirement for antimicrobials and ensure prudent use. Dairy Australia's Countdown program (Brightling et al, 2009; <https://www.dairyaustralia.com.au/farm/animal-management/mastitis/countdown>), focuses on reducing mastitis and improving milk quality, and has been active since 1998. A flagship project within Countdown is the adviser training short course, which has now evolved into year-long Countdown Milk Quality Advisor training. These upskill all dairy advisors (veterinarians, field services and milking machine technicians) in milk quality investigations and on farm communication to change practises on farm, including promoting prudent antimicrobial use. The development of the dry cow consult tool (<http://drycowconsult.com.au/>; through the Dairy Australia Countdown project) assists farmers (in conjunction with their veterinarian) to adopt selective treatment of cows with antibiotic therapy through the dry period, rather than routine herd treatment.

Most of Dairy Australia's other animal health extension programs arguably contribute materially to the judicious use of antimicrobials, the details of which can be found on the Dairy Australia website (<https://www.dairyaustralia.com.au/>). The history of key AMS initiatives in the Australian dairy industry are outlined in Appendix 4. The key messages from many of these programs focus on disease prevention and consequent optimisation of antimicrobial use to reduce the risk of AMR bacteria occurring. These programs improve cow management and reduce reliance on feed additives which alter ruminal fermentation (i.e. microbial growth).

The Healthy Hooves program (more information: <http://healthyhooves.dairyaustralia.com.au/>) aims to reduce lameness in herds, which is something that may need to be treated with antibiotics. In the area of calf management, the Rearing Healthy Calves program (more information: <https://www.dairyaustralia.com.au/healthycalves>) has a strong focus on highlighting judicious use of antimicrobials and as well as the importance of improved colostrum practise, which can reduce the risk of infections in calves by improving immunity. Implementation of good biosecurity is paramount to disease prevention.

Dairy Australia, along with state government stakeholders, is in the process of developing a biosecurity tool for dairy farmers. The ACV (a special interest group of the AVA) have an online biosecurity program, called BIOCHECK® (<https://www.ava.com.au/cattle/biocheck-public>) which is a program designed to improve farm preparedness for major biosecurity risks and ensure appropriate risk management strategies are in place on farm.

Involvement of the Australian Dairy industry in Australian livestock antimicrobial stewardship

The Australian dairy industry has been contributing to collective Australian antimicrobial stewardship efforts since they were formalised in 2014/15 through the National AMR strategy (Australian Government, 2015). This involvement has led to industry-level initiatives aimed at better understanding antimicrobial usage on Australian dairy farms through two surveys (2012-13 and 2016-17) of veterinary antibiotic sales data based on aggregated quantity (kg) of actives sold.

The dairy industry is in the process of developing a comprehensive AMS strategy to improve adoption of AMS and judicious use principles. This includes improved understanding of what constitutes best practice management of antibiotics in dairy farming without compromising the health and welfare of Australia's dairy herd, monitoring antimicrobial consumption, implementation of initiatives to improve adoption of judicious use principles and good antimicrobial stewardship practices. A strategy for AMS in the dairy industry is being developed for inclusion in the Australian Dairy Industry Sustainability Framework (<http://www.sustainableairyoz.com.au/>). Whilst still in the early stages, the draft principles of AMS in the dairy industry are coordinated alongside the national livestock AMS approach and cover advancing and refining efforts to implement AMS principles, prioritisation of activities based on risk assessment and risk management, disease prevention and refinement of antimicrobial use and shared responsibility through chain.



CONCLUSION

This report cannot be considered comprehensive, as there is an abundance of resources available online and elsewhere that support the health, welfare, husbandry (including hygiene) and disease prevention in livestock through biosecurity. However, the purpose of this report is to provide an overview of the historical and current practises relevant to AMS in each of the contributing industries that are already in place and the information that is available to support AMS in Australian livestock. In particular, this report is directed at the stakeholders that currently lack a comprehensive understanding of how livestock industries operate in Australia. The implementation of the National AMR Strategy (Australian Government, 2015), and the process for developing and implementing the next national strategy, have provided an impetus for each industry to assess its current activities under an AMS framework and identify opportunities for improvements and define barriers that align across all animal industries which can be addressed collaboratively. The National AMR Strategy also helps to identify pathways for collaboration between governments and industries to consolidate resources in an effort to address some of these barriers. For example, the processes that were coordinated through the Office of the Australian Chief Veterinary Office in 2017-18, with more planned in the future, and the coordination of an Australian conference specifically on AMS to be held November 11-13th 2018. This conference will help stakeholders in the Australian animal industries, including the livestock industries, with their understanding of AMS and its principles, and highlight barriers and improvements required for successful AMS programmes.

It is expected that this report will be regularly updated to capture information from additional industries, and updates on the initiatives from industries that were in a position to contribute to this first version.

While AMR is a significant human health issue, it should not be forgotten that it is also an animal health issue, as antimicrobials (including those not used in human medicine) are still required to manage disease in livestock, and animal pathogens with resistance to these antimicrobials can potentially cause significant issues to both humans and livestock. Ignoring AMR in animals may also result in the use of antimicrobials of importance, or of greater importance, to human medicine, which could perpetuate the development of further AMR. Despite this, the momentum is high in Australia, 2018, for adopting AMS principles, through capturing current efforts and identifying further areas for improvement. Continuation of this momentum through collaborative efforts will

be essential to maintaining and improving Australia's positive track record for AMS, and generally low levels of AMR bacteria in livestock. The development and implementation of systems to capture the information needed to support this narrative at the national level is essential to minimising AMR into the future.

ACKNOWLEDGEMENTS

The author would like to acknowledge the substantial contributions, direction and review provided by those that contributed to this report from the AVA, state government, independent experts and the red meat, poultry, pork and dairy industries. Due acknowledgement also goes to the Australian Government Department of Agriculture and Water Resources for not only providing the funding for the development of this report, through the Australian Government's *Animal Biosecurity and Response Reform* program, but also for the input and direction throughout.



REFERENCES

- Abraham, S., O'Dea, M., Page, S., and Trott, D. (2017). Current and future antimicrobial resistance issues for the Australian pig industry. *Animal Production Science* 57, 2398-2407.
- ACMF (2018a). "Surveillance for antimicrobial resistance in enteric commensals and pathogens in Australian meat chickens". Australian Chicken Meat Federation, North Sydney, New South Wales.
- ACMF (2018b). "Responsible Use of Antibiotics in the Australian Chicken Meat Industry." Australian Chicken Meat Federation, North Sydney, New South Wales.
- AHA (2018). "Animal Health in Australia; 2017." Animal Health Australia, Canberra, Australia.
- APL (2013a). "Care of the Compromised Pig." Australian Pork Limited, Australian Capital Territory.
- APL (2013b). "Companion Handbook to the Model Code of Practice for the Welfare of Pigs (3rd ed.)." Australian Pork Limited, Australian Capital Territory.
- APVMA (2014). "Report on the Quantity of Antimicrobial Products Sold for Veterinary Use in Australia: July 2005 to June 2010 ". Australian Pesticides and Veterinary Medicines Authority, The Australian Government, Canberra.
- APVMA (2015). "Using veterinary medicines." Australian Pesticides and Veterinary Medicines Authority, The Australian Government, Canberra.
- APVMA (2017). "Antibiotic resistance in animals." Australian Pesticides and Veterinary Medicines Authority, The Australian Government, Canberra.
- APVMA (2018). "APVMA Legislative Framework." Australian Pesticides and Veterinary Medicines Authority, The Australian Government, Canberra.
- ARSC (2014). "Importance Ratings and Summary of Antibacterial Uses in Humans in Australia." Antimicrobial Resistance Standing Committee, The Australian Government, Canberra.
- Australian Government (1961). "National Residue Survey." Department of Agriculture and Water Resources, Canberra, Australia.
- Australian Government (2007). "Pilot Surveillance Program for Antimicrobial Resistance in Bacteria of Animal Origin." Department of Agriculture and Water Resources, Canberra, Australia.
- Australian Government (2015). "Responding to the threat of antimicrobial resistance: Australia's First National Antimicrobial Resistance Strategy 2015–2019." Department of Health and Department of Agriculture, Canberra, ACT.
- Australian Government (2016). "Implementation Plan: Australia's First National Antimicrobial Resistance Strategy ". Department of Health and Department of Agriculture, Canberra, Australia.
- APIA (1987). "The Use of Antibiotics and Other Drugs in the Poultry Industry - A Code of Practice for Use by Companies". Australian Poultry Industries Association, North Sydney, NSW, Australia.
- APV (1989). "Code of Practice for Use of Veterinary Medicines in the Pig Industry." Australian Pig Veterinarians.
- ASTAG (Australian Strategic and Technical Advisory Group on AMR) (2018). "Importance Ratings and Summary of Antibacterial Uses in Human and Animal Health in Australia, Version 1.0". The Australian Government, Canberra.
- AVA (2013). "Guidelines for prescribing, authorising and dispensing veterinary medicines". Australian Veterinary Association, St Leonards, NSW.
- AVA (2017). "Antibiotic prescribing guidelines for livestock and horses ". Australian Veterinary Association, St Leonards, NSW.
- AVA (2018). "The Australian Veterinary Association's (AVA) strategic priority program." Australian Veterinary Association, St Leonards, NSW.
- AVPA (2001). "AVPA Code of Practice for the Use of Antibiotics in the Poultry Industry – 2001 Edition." Australasian Veterinary Poultry Association.
- Bagust, T. J. (1989). "An overview of Australia's poultry industry in 1989." *Australian Veterinary Journal* 66(12): 416-418.
- Barlow, R. S., McMILLAN, K. E., Duffy, L. L., Fegan, N., Jordan, D., and Mellor, G. E. (2015). Prevalence and antimicrobial resistance of *Salmonella* and *Escherichia coli* from Australian cattle populations at slaughter. *Journal of food protection* 78, 912-920.
- Barlow, R. S., McMillan, K. E., Duffy, L. L., Fegan, N., Jordan, D., and Mellor, G. E. (2017). Antimicrobial resistance status of *Enterococcus* from Australian cattle populations at slaughter. *PLoS one* 12, e0177728.
- Barnes, T. S., Hay, K., Morton, J. M. M and Mahony, T. J (2015). Epidemiology and management of bovine respiratory disease in feedlot cattle.
- Barton M and Wilkins J, R. (2001). Antibiotic resistance in bacteria isolated from poultry.
- Brightling, P, B., Dyson, R. D., Hope, A. F. and Penry, J. (2009). A national programme for mastitis control in Australia: Countdown Downunder. *Irish Veterinary Journal* 62 (Suppl 4), S52-S58.
- Clements, M. (2015). Hemorrhagic enteritis vaccine for Australian turkey sector. WATTAGNet article: <https://www.wattagnet.com/articles/25238-hemorrhagic-enteritis-vaccine-for-australian-turkey-sector>
- Colditz, I. G., and Hine, B. C. (2016). Resilience in farm animals: biology, management, breeding and implications for animal welfare. *Animal Production Science* 56, 1961-1983.
- Collineau, L., Belloc, C., Stärk, K. D., Hémonic, A., Postma, M., Dewulf, J., and Chauvin, C. (2017). Guidance on the selection of appropriate indicators for

- quantification of antimicrobial usage in humans and animals. *Zoonoses and public health* 64, 165-184.
- Cudmore JW, W. P., McGahan EJ and Sullivan TJ (2015). "NFAS Review." Meat & Livestock Australia, Sydney, New South Wales.
- Cunha, C. B. (2018). *Antimicrobial Stewardship Programs: Principles and Practice*. Medical Clinics of North America.
- Dairy Australia (2018). "Annual Report 2017-18". Dairy Australia, Southbank, Victoria.
- Darden, H. (1988). *The Stewardship of Planet Earth: Preserving Biological Diversity*. Bios, 99-103.
- Delaplane, J., and Stuart, H. (1943). The propagation of a virus in embryonated chicken eggs causing a chronic respiratory disease of chickens. *Am. J. Vet. Res* 4, 325-332.
- EMA and ESVAC (2017). "Sales of veterinary antimicrobial agents in 30 European countries in 2015; Trends from 2010 to 2015."
- Gaskins, H., Collier, C., and Anderson, D. (2002). Antibiotics as growth promotants: mode of action. *Animal biotechnology* 13, 29-42.
- Guardabassi, L., and Prescott, J. F. (2015). Antimicrobial stewardship in small animal veterinary practice: from theory to practice. *Veterinary Clinics: Small Animal Practice* 45, 361-376.
- Harbarth, S., and Hackett, J. (2018). Introduction: DRIVE-AB's definitions and indicators to monitor responsible antibiotic use. *Journal of Antimicrobial Chemotherapy* 73, vi2-vi2.
- Hardefeldt, L., Marena, M., Crabb, H., Stevenson, M., Gilkerson, J., Billman-Jacobe, H., and Browning, G. (2018). Antimicrobial susceptibility testing by Australian veterinary diagnostic laboratories. *Australian veterinary journal* 96, 142-146.
- Hassan, Y. I., Lahaye, L., Gong, M. M., Peng, J., Gong, J., Liu, S., Gay, C. G., and Yang, C. (2018). Innovative drugs, chemicals, and enzymes within the animal production chain. *Veterinary research* 49, 71.
- Hermesch, S., and Dominik, S. (2014). *Breeding Focus 2014-improving resilience*. Animal Genetics and Breeding Unit, University of New England: Armidale.
- Hoelzer, K., Bielke, L., Blake, D. P., Cox, E., Cutting, S. M., Devriendt, B., Erlacher-Vindel, E., Goossens, E., Karaca, K., and Lemiere, S. (2018a). Vaccines as alternatives to antibiotics for food producing animals. Part 1: challenges and needs. *Veterinary research* 49, 64.
- Hoelzer, K., Bielke, L., Blake, D. P., Cox, E., Cutting, S. M., Devriendt, B., Erlacher-Vindel, E., Goossens, E., Karaca, K., and Lemiere, S. (2018b). Vaccines as alternatives to antibiotics for food producing animals. Part 2: new approaches and potential solutions. *Veterinary research* 49, 70.
- Holloway, S. (2013). *Antibiotic Prescribing Detailed Guidelines: Australasian Infectious Diseases Advisory Panel*.
- Jee, Y., Carlson, J., Rafai, E., Musonda, K., Huong, T. T. G., Daza, P., Sattayawuthipong, W., and Yoon, T. (2018). Antimicrobial resistance: a threat to global health. *The Lancet Infectious Diseases* 18, 939-940.
- JETACAR (1999). "The use of antibiotics in food-producing animals: antibiotic-resistant bacteria in animals and humans," Rep. No. 1864960612. Joint Expert Technical Advisory Committee on Antibiotic Resistance, The Australian Government Commonwealth Department of Health and Aged Care and Commonwealth Department of Agriculture, Fisheries and Forestry.
- Jordan, D., Chin, J. C., Fahy, V., Barton, M. D., Smith, M., and Trott, D. (2009). Antimicrobial use in the Australian pig industry: results of a national survey. *Australian Veterinary Journal* 87, 222-229.
- Kidsley, A. K., Abraham, S., Bell, J. M., O'Dea, M., Laird, T. J., Jordan, D., Mitchell, P., McDevitt, C. A., and Trott, D. J. (2018). Antimicrobial susceptibility of *Escherichia coli* and *Salmonella* spp. isolates from healthy pigs in Australia: results of a pilot national survey. *Frontiers in microbiology* 9.
- Le Maréchal, M., Tebano, G., Monnier, A. A., Adriaenssens, N., Gyssens, I. C., Huttner, B., Milanič, R., Schouten, J., Stanić Benić, M., and Versporten, A. (2018). Quality indicators assessing antibiotic use in the outpatient setting: a systematic review followed by an international multidisciplinary consensus procedure. *Journal of Antimicrobial Chemotherapy* 73, vi40-vi49.
- Lillehoj, H., Liu, Y., Calsamiglia, S., Fernandez-Miyakawa, M. E., Chi, F., Cravens, R. L., Oh, S., and Gay, C. G. (2018). Phytochemicals as antibiotic alternatives to promote growth and enhance host health. *Veterinary research* 49, 76.
- Lloyd, D. H., and Page, S. W. (2018). *Antimicrobial Stewardship in Veterinary Medicine*. Microbiology spectrum 6.
- Mifflin J and Blackall P, R. (1998). Development of diagnostic tools for the rapid and specific detection of avian bacterial pathogens.
- MLA (2014). "Antimicrobials and the cattle industry." Meat & Livestock Australia, Sydney, New South Wales.
- MLA (2017). "State of the Industry Report: The Australian Red Meat and Livestock Industry." Meat & Livestock Australia, Sydney, New South Wales.
- MLA (2018a). "Antimicrobial stewardship guidelines for the Australian cattle feedlot industry." Meat & Livestock Australia, Sydney, New South Wales.
- MLA (2018b). "National Feedlot Accreditation Scheme Rules". Meat & Livestock Australia, Sydney, New South Wales.
- Neethirajan, S. (2017). Recent advances in wearable sensors for animal health management. *Sensing and Bio-Sensing Research* 12, 15-29.
- Obeng, A., Rickard, H., Sexton, M., Pang, Y., Peng, H., and Barton, M. (2012a). Antimicrobial susceptibilities and resistance genes in *Campylobacter* strains isolated from poultry and pigs in Australia. *Journal of applied microbiology* 113, 294-307.
- Obeng, A. S., Rickard, H., Ndi, O., Sexton, M., and Barton, M. (2012b). Antibiotic resistance, phylogenetic grouping and virulence potential of *Escherichia coli* isolated from the faeces of intensively farmed and free range poultry. *Veterinary microbiology* 154, 305-315.
- Obeng, A. S., Rickard, H., Ndi, O., Sexton, M., and Barton, M. (2014). Prevalence of Antimicrobial Resistance in Enterococci and *Escherichia coli* in Meat

- Chicken Flocks During a Production Cycle and Egg Layer Pullets During Rearing. *International Journal of Poultry Science* 13, 489.
- Page, S., Prescott, J., and Weese, S. (2014). The 5Rs approach to antimicrobial stewardship. *Veterinary Record* 175, 207-208.
- Pande, V. V., Gole, V. C., McWhorter, A. R., Abraham, S., and Chousalkar, K. K. (2015). Antimicrobial resistance of non-typhoidal *Salmonella* isolates from egg layer flocks and egg shells. *International journal of food microbiology* 203, 23-26.
- PISC (2002). "Model code of practice for the welfare of animals: Domestic poultry," Rep. No. 0643068678. Primary Industries Standing Committee, CSIRO Publishing, Australia.
- Prescott, J. F., and Boerlin, P. (2016). Antimicrobial use in companion animals and Good Stewardship Practice. *The Veterinary Record* 179, 486.
- Richeson, J. T., Lawrence, T. E., and White, B. J. (2018). Using advanced technologies to quantify beef cattle behavior. *Translational Animal Science* 2, 223-229.
- Scott Weese, J., Page, S. W., and Prescott, J. F. (2013). Antimicrobial stewardship in animals. *Antimicrobial therapy in veterinary medicine*, 117-132.
- Seal, B. S., Drider, D., Oakley, B. B., Brüssow, H., Bikard, D., Rich, J. O., Miller, S., Devillard, E., Kwan, J., and Bertin, G. (2018). Microbial-derived products as potential new antimicrobials. *Veterinary research* 49, 66.
- Shaban, R., Simon, G., Trott, D., Turnidge, J., and Jordan, D. (2014). Surveillance and reporting of antimicrobial resistance and antibiotic usage in animals and agriculture in Australia. Department of Agriculture, The Australian Government: Canberra.
- Simjee, S., McDermott, P., Trott, D., and Chuanchuen, R. (2018). Present and Future Surveillance of Antimicrobial Resistance in Animals: Principles and Practices. *Microbiology spectrum* 6.
- Smith, D. R., Dolk, F. C. K., Pouwels, K. B., Christie, M., Robotham, J. V., and Smieszek, T. (2018). Defining the appropriateness and inappropriateness of antibiotic prescribing in primary care. *Journal of Antimicrobial Chemotherapy* 73, ii11-ii18.
- van Dijk, A., Hedegaard, C. J., Haagsman, H. P., and Heegaard, P. M. (2018). The potential for immunoglobulins and host defense peptides (HDPs) to reduce the use of antibiotics in animal production. *Veterinary research* 49, 68.
- Van, T. T. H., Elshagmani, E., Gor, M. C., Scott, P. C., and Moore, R. J. (2016). *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *International journal of systematic and evolutionary microbiology* 66, 4518-4524.
- Van, T. T. H., Gor, M.-C., Anwar, A., Scott, P. C., and Moore, R. J. (2017). *Campylobacter hepaticus*, the cause of spotty liver disease in chickens, is present throughout the small intestine and caeca of infected birds. *Veterinary microbiology* 207, 226-230.
- Versporten, A., Gyssens, I. C., Pulcini, C., Monnier, A. A., Schouten, J., Milanič, R., Stanić Beničić, M., Tebano, G., Le Maréchal, M., and Zanichelli, V. (2018). Metrics to assess the quantity of antibiotic use in the outpatient setting: a systematic review followed by an international multidisciplinary consensus procedure. *Journal of Antimicrobial Chemotherapy* 73, vi59-vi66.
- Whithear, K., Harrigan, K., and Ghiocas, E. (1990). Safety of temperature sensitive mutant *Mycoplasma gallisepticum* vaccine. *Australian veterinary journal* 67, 159-165.
- Whithear, K., Htwe, T., and Sulaiman, I. (1986). Survey of disc diffusion antimicrobial sensitivity testing in avian bacteriology laboratories and the evaluation of a standardised method. *Australian veterinary journal* 63, 113-117.

APPENDIX 1

References for the Veterinary AMS Timeline

- AIDAP (2013). Antibiotic Prescribing Detailed Guidelines, Australasian Infectious Diseases Advisory Panel.
- AIDAP (2016). Practical infection control guidelines, Australasian Infectious Diseases Advisory Panel.
- Australian Government (2015). National Antimicrobial Resistance Strategy 2015–2019. Canberra, ACT, Department of Health and Department of Agriculture.
- APIA (1987). "The Use of Antibiotics and Other Drugs in the Poultry Industry - A Code of Practice for Use by Companies". Australian Poultry Industries Association, North Sydney, NSW.
- AVA (2011). Australian Veterinary Association Guidelines for Veterinary Personal Biosecurity Version 1.0. St Leonards, NSW, Australian Veterinary Association.
- AVA (2013). Australian Veterinary Association Guidelines for Veterinary Personal Biosecurity 2nd edition. St Leonards, NSW, Australian Veterinary Association.
- AVA (2017). Australian Veterinary Association Guidelines for Veterinary Personal Biosecurity 3rd edition. St Leonards, NSW, Australian Veterinary Association.
- AVA (Australian Veterinary Association) (2005). Guidelines for Prescribing, Authorising and Dispensing Veterinary Medicines. M. P. Bond. Canberra, Australian Veterinary Association.
- Barlow, R. and K. Gobius (2008). Pilot survey for antimicrobial resistant (AMR) bacteria in Australian food. Cannon Hill, Queensland, Food Science Australia.
- Barton, M. (2017). Antibiotic Resistance in Animals - a Report for the APVMA. Canberra, Australian Pesticides and Veterinary Medicines Authority (APVMA).
- Cheng, A. C., J. Turnidge, P. Collignon, D. Looke, M. Barton and T. Gottlieb (2012). "Control of fluoroquinolone resistance through successful regulation, Australia." *Emerg Infect Dis* 18(9): 1453-1460.
- Commonwealth Department of Health and Aged Care and Commonwealth Department of Agriculture; Fisheries and Forestry (2000). The Commonwealth Government Response to the Report of the Joint Expert Technical Advisory Committee on Antibiotic Resistance (JETACAR). Canberra, Commonwealth of Australia.
- Commonwealth Interdepartmental JETACAR Implementation Group (CIJIG) (2002). Progress Report. Canberra, Commonwealth Department of Health & Aged Care.
- Cooper (Editor), B. S. (1994). Antimicrobial prescribing guidelines for veterinarians. Sydney, NSW, University of Sydney, Post Graduate Foundation in Veterinary Science in association with National Health & Medical Research Council.
- Cooper (Editor), B. S. (2000). Antimicrobial prescribing guidelines for veterinarians (2nd edition). Sydney, NSW, University of Sydney, Post Graduate Foundation in Veterinary Science in association with National Health & Medical Research Council.
- DAFF (2007). Pilot Surveillance Program for Antimicrobial Resistance in Bacteria of Animal Origin. November 2003 to July 2004. Canberra, ACT, Australian Government Department of Agriculture, Fisheries and Forestry (DAFF).
- Doyle, K. (2001). JETACAR - The Role of Professional Bodies. National Summit on Antibiotic Resistance Stamford Sydney Airport Hotel, Commonwealth Department of Health and Aged Care and the Commonwealth Department of Agriculture, Fisheries and Forestry.
- Gilbert, G. L. (1994). Foreward. Antimicrobial prescribing guidelines for veterinarians. B. S. Cooper. Sydney, NSW, University of Sydney, Post Graduate Foundation in Veterinary Science in association with National Health & Medical Research Council.
- Gottlieb, T. and G. R. Nimmo (2011). "Antibiotic resistance is an emerging threat to public health: an urgent call to action at the Antimicrobial Resistance Summit 2011." *Medical Journal of Australia* 194(6): 281-283.
- Joint Expert Technical Advisory Committee on Antibiotic Resistance (JETACAR) (1999). The Use of Antibiotics in Food-Producing Animals: Antibiotic-Resistant Bacteria in Animals and Humans. Canberra, Commonwealth Department of Health and Aged Care and Commonwealth Department of Agriculture, Fisheries and Forestry: i-xxxv, 1-224.
- Jones, C. and B. Howden (2017). Report from the 2nd National AMR Summit, June 29, 2017. <https://www.asid.net.au/documents/item/1423>, Australasian Society for Infectious Diseases (ASID) and Australian Society for Antimicrobials (ASA).
- McEwen, J. (2007). A History of Therapeutic Goods Regulation in Australia. Canberra, Commonwealth of Australia.
- National Drugs and Poisons Schedule Committee (NDPSC) (2003). Record of Reasons 39th Meeting 14-16 October 2003. Antibiotics for consideration following recommendations of the Joint Expert Technical Advisory Committee on Antibiotic Resistance (JETACAR). Canberra, Australian Government Department of Health and Ageing.
- National Registration Authority (NRA) (2000). Part 10 of Veterinary Requirement Series. Submission to the Working Party on Antibiotics in support of the approval of new antibiotic active constituents and registration of veterinary chemical products containing antibiotic active constituents, and significant variations to such products (new dosage forms, extensions to use). Kingston, ACT, National Registration Authority.
- NHMRC (1977). Report of the Eighty-Third Session, Hobart, April 1977. Canberra, Commonwealth Department of Health, National Health and Medical Research Council.

- Page, S. W. (2000). The future of veterinary antibiotics. AVA Annual Conference. L. Cook. Perth, Western Australia, Australian Veterinarians in Industry and Australian Veterinarians in Public Health: 43-46.
- Ryan, S., H. Polley, R. Di Natale, A. McEwen, A. Sinodinos, J. Williams and N. Xenophon (2013). Progress in the implementation of the recommendations of the 1999 Joint Expert Technical Advisory Committee on Antibiotic Resistance. Parliament House, Canberra, The Senate Finance and Public Administration References Committee.
- Therapeutic Goods Administration (TGA) (2013). Final decisions & reasons for decisions by delegates of the Secretary to the Department of Health November 2013. Part B - Final decisions on matters not referred to an expert advisory committee. 4. Chemicals. 4.1 Tylosin (implementation date of 1 June 2014). Canberra, Australian Government Department of Health.
- Tolhurst, J. C., G. Buckle and S. W. Williams (1955). Chemotherapy with Antibiotics and Allied Drugs. Canberra, National Health and Medical Research Council.
- Webber, J. (2006). A comprehensive integrated surveillance program to improve Australia's response to antimicrobial resistance (August 2006). A report prepared for the NHMRC's Expert Advisory Group on Antimicrobial Resistance (EAGAR). Canberra, EAGAR.

APPENDIX 2

Antibacterial agents registered for antibacterial use in livestock by APVMA (information accessed October 2018)

Antibacterial Agent	Class	Importance ^a		Cattle	Sheep	Pig	Meat Chicken	Egg-layer hen
		ASTAG 2018	WHO 2017					
Novobiocin	Aminocoumarin	nhu	5	X ^o				
Spectinomycin	Aminocyclitol	med	4			X	X	X ^{e,r}
Apramycin	Aminoglycoside	med	2	X		X	X	
Dihydrostreptomycin	Aminoglycoside	low	2	X				
Framycetin	Aminoglycoside	low	2	X	X			
Neomycin	Aminoglycoside	low	2	X	X	X	X	X ^{e,r}
Streptomycin	Aminoglycoside	low	2	X				
Trimethoprim ^s	Diaminopyrimidine	med	3	X	X	X	X	X ^r
Flavophospholipol	Glycophospholipid	nhu	5	X		X	X	X ^{e,r}
Lasalocid	Ionophore	nhu	5	X			X ^f	X ^r
Monensin	Ionophore	nhu	5	X	X		X ^f	X ^r
Narasin	Ionophore	nhu	5	X			X ^f	
Salinomycin	Ionophore	nhu	5	X		X	X ^f	X ^r
Lincomycin	Lincosamide	med	3	X		X	X	X ^{e,r}
Erythromycin	Macrolide	low	1	X	X	X	X	
Oleandomycin	Macrolide	low	1	X ^o				
Tilmicosin	Macrolide	low	1	X		X		
Tulathromycin	Macrolide	low	1	X		X		
Tylosin	Macrolide	low	1	X		X	X	X ^r
Avilamycin	Orthosomycin	nhu	5				X	
Florfenicol	Phenicol	low	3	X		X		
Tiamulin	Pleuromutilin	mhu	4			X	X	
Bacitracin	Polypeptide i	low	4	X	X		X	X ^{e,r}
Polymyxin B	Polypeptide ii	high	1	X	X			
Olaquinox	Quinoxaline	nhu	5			X		
Virginiamycin	Streptogramin	high	3	X	X		X	
SulfadiazineT ⁺	Sulfonamide	low	3	X	X	X	X	
SulfadimidineT ^{+/-}	Sulfonamide	low	3	X	X	X	X	X ^r
SulfadoxineT ⁺	Sulfonamide	low	3	X	X	X		
Chlortetracycline	Tetracycline	low	3	X		X	X	X ^{e,r}
Oxytetracycline	Tetracycline	low	3	X	X	X	X	
Cephapirin	β lactam cephalosporin [1GC]	med	3	X				
Cephalonium	β lactam cephalosporin [1GC]	med	3	X ^o				
Cefuroxime	β lactam cephalosporin [2GC]	med	3	X ^o				
Ceftiofur	β lactam cephalosporin [3GC]	high	1	X				
Amoxicillin	β lactam penicillin	low	2	X	X	X	X	X ^r
Ampicillin	β lactam penicillin	low	2	X ^o				
Cloxacillin	β lactam penicillin	med	3	X	X			
Penethamate	β lactam penicillin	low	2	X	X	X		
Penicillin (and salts)	β lactamase inhibitor	low	2	X	X	X		
Clavulanic acid		med	2	X ^o				

^a IMPORTANCE for human medicine: ASTAG, version 1.0 2018; mhu minor human use; nhu no human use; WHO, version 5 2017; 1 HPCIA; 2 CIA; 3 HIA; 4 IA; 5 nhu, ^s combination with a sulfonamide; T+/- with or without trimethoprim, ^o active only available in an intramammary product, ^e discard eggs for 7 days; ^r pullet laying replacement, ^f Label claim for coccidiosis, not antibacterial activity

Only antibacterial agents contained within products with antibacterial claims are included in this list. Other active constituents with antibacterial activity with claims for other uses are not included, for example, maduramicin, semduramicin and sulfaquinoxaline. Different products may have different label directions.

APPENDIX 3

Vaccines registered, or approved by permit, for use in cattle, pigs and poultry by APVMA (information accessed October 2018)

<i>Immunogen</i>	<i>Type</i>	<i>Cattle</i>	<i>Sheep</i>	<i>Pig</i>	<i>Poultry</i>
<i>Actinobacillus pleuropneumoniae</i>	Bacteria			X	
<i>Anaplasma centrale</i>	Protozoa	X			
Avian encephalomyelitis virus	Virus				X
Avian influenza virus (H5N2, H5N9, H7N1)	Virus				X
<i>Avibacterium paragallinarum</i>	Bacteria				X ^a
<i>Babesia bigemina</i>	Protozoa	X			
<i>Babesia bovis</i>	Protozoa	X			
<i>Bacillus anthracis</i> (Sterne 34F2 strain)	Bacteria	X	X	X	
Bovine coronavirus	Virus	X			
Bovine ephemeral fever virus	Virus	X			
Bovine herpesvirus 1	Virus	X			
Bovine pestivirus	Virus	X			
Bovine rotavirus	Virus	X			
<i>Campylobacter fetus</i> subsp <i>fetus</i>	Bacteria	X	X		
<i>Campylobacter fetus</i> subsp <i>venerealis</i>	Bacteria	X			
<i>Campylobacter hepaticus</i>	Bacteria				X ^a
<i>Campylobacter jejuni</i> subsp <i>jejuni</i>	Bacteria		X		
Chicken anaemia virus	Virus				X
<i>Clostridium botulinum</i> Type C	Bacteria	X	X		
<i>Clostridium botulinum</i> Type D	Bacteria	X	X		
<i>Clostridium chauvoei</i>	Bacteria	X	X		
<i>Clostridium haemolyticum</i>	Bacteria	X	X		
<i>Clostridium novyi</i> Type B	Bacteria	X	X		
<i>Clostridium perfringens</i> Type A	Bacteria				X ^a
<i>Clostridium perfringens</i> Type B	Bacteria	X	X		
<i>Clostridium perfringens</i> Type C	Bacteria	X	X		
<i>Clostridium perfringens</i> Type D	Bacteria	X	X		
<i>Clostridium septicum</i>	Bacteria	X	X		
<i>Clostridium tetani</i>	Bacteria	X	X		
Contagious pustular dermatitis virus (Orf virus)	Virus		X		
<i>Corynebacterium pseudotuberculosis</i>	Bacteria		X		
<i>Dichelobacter nodosus</i>	Bacteria		X		
Egg drop syndrome 76 virus	Virus				X
<i>Eimeria acervulina</i>	Protozoa				X
<i>Eimeria brunetti</i>	Protozoa				X
<i>Eimeria maxima</i>	Protozoa				X
<i>Eimeria mitis</i>	Protozoa				X
<i>Eimeria necatrix</i>	Protozoa				X
<i>Eimeria praecox</i>	Protozoa				X
<i>Eimeria tenella</i>	Protozoa				X
<i>Erysipelothrix rhusiopathiae</i>	Bacteria		X ^a	X	X

Escherichia coli (many types)	Bacteria	X		X	X
Foot and mouth disease virus (FMD)	Virus	X ^a	X ^a	X	
Fowl adenovirus	Virus				X
Fowl pox virus	Virus				X
Haemophilus parasuis	Bacteria			X	
Herpes virus of turkeys (HVT)	Virus				X
Infectious bronchitis virus (IB)	Virus				X
Infectious bursal disease virus (IBD)	Virus				X
Infectious laryngotracheitis virus (ILT)	Virus				X
Klebsiella pneumoniae	Bacteria			X ^a	
Lawsonia intracellularis	Bacteria			X	
Leptospira borgpetersenii serovar Hardjo	Bacteria	X	X		
Leptospira interrogans serovar Pomona	Bacteria	X	X	X	
Leptospira interrogans serovar Tarassovi	Bacteria			X	
Mannheimia haemolytica	Bacteria	X			
Marek's disease virus	Virus				X
Moraxella bovis	Bacteria	X			
Mycobacterium paratuberculosis	Bacteria	X	X		
Mycoplasma gallisepticum	Bacteria				X
Mycoplasma hyopneumoniae	Bacteria			X	
Mycoplasma synoviae	Bacteria				X
Newcastle disease virus	Virus				X
Ornithobacterium rhinotracheale (ORT)	Bacteria				X ^a
Papilloma virus	Virus	X ^a	X ^a	X ^a	
Pasteurella multocida	Bacteria	X ^a	X ^a	X	X
Porcine circovirus type 2	Virus			X	
Porcine parvovirus	Virus			X	
Salmonella Bovismorbificans, Uganda & Zanzibar	Bacteria	X			
Salmonella Dublin	Bacteria	X			
Salmonella Typhimurium	Bacteria	X			X
Staphylococcus hyicus	Bacteria			X ^a	
Staphylococcus spp (incl aureus, epidermidis, hyicus)	Bacteria	X ^a	X ^a	X ^a	X ^a
Streptococcus suis	Bacteria			X ^a	

^a vaccine only available under permit, APVMA PubCRIS and Permit databases accessed 11 October 2018

APPENDIX 4

Historical Antimicrobial Stewardship efforts in the Australian Dairy Industry.

<i>Timeline</i>	<i>Event</i>	<i>Reference available from</i>
1990s (& ongoing)	Dairy processor engagement in monitoring antibiotic use and residue risk management (Rapid Screen tests/monitoring)	Dairy Australia
1996 (& ongoing)	Australian Milk Residue Analysis survey is undertaken annually to provide assurance about milk product safety and integrity, highlighting the effectiveness of dairy industry biosecurity measures and management systems, including controls on veterinary products. EU requirements have been a key influence on the design of the survey. Furthermore: <ul style="list-style-type: none"> • The National Residue Survey has the same assurance function for meat and meat products • The National Antibacterial Residue Minimisation Program minimises the risk of violative antibacterial residues in bobby calves 	https://www.dairyaustralia.com.au/industry/food-safety-and-regulation/regulatory-framework/australian-milk-residue-analysis-survey
1998 (& ongoing)	Dairy Australia's AMS flagship project Countdown: to assist farmers in mastitis management and prudent antimicrobial use	https://www.dairyaustralia.com.au/countdown
2000 (& ongoing)	Australian Milk Quality Awards – annual dairy farmer award based on Bulk Milk Cell Count – a key indicator of good mastitis control and public acknowledgement of individual farmers.	https://www.dairyaustralia.com.au/farm/animal-management/mastitis/milk-quality-awards/2017-milk-quality-awards
2006 (& ongoing)	Cups On Cups Off milking training program developed by Dairy Australia under the Countdown program for milkers and lactation managers to prevent and control mastitis and manage milk quality in the dairy	Dairy Australia
2008	Primary Production and Processing Standards for Dairy Products (4.2.4 Food Standards Code), which mandates on-farm food safety programs subject to regular audit by state food safety regulators, including appropriate use of veterinary medicines.	https://www.legislation.gov.au/Series/F2012L00294
2009	Zoetis Survey of AMR in mastitis pathogens	Dairy Australia
2010 (& ongoing)	Dairy Australia, Australian Dairy Farmers and Victorian Government joint Vet Investigation Residue Management Project to minimize AM residues in dairy calf products	Dairy Australia
2011 (& ongoing)	Dairy Australia Healthy Calves Project to improve calf health and reduce reliance on antibiotics	https://www.dairyaustralia.com.au/healthycalves
2013	IDF release of Guide to Prudent Use of Antimicrobial Agents in Dairy Production - International Working Group led by Dairy Australia	Dairy Australia
2013	First survey of antibiotic use in Australian dairy industry by dairy cattle veterinarians (AMU from vet sales data)	Dairy Australia
2014	Survey of AMR in enteric organisms from dairy, beef and feedlot cattle	Dairy Australia
2014	Countdown 2020: Farm Guidelines for Mastitis Control	Dairy Australia
2014	Countdown Mastitis Toolkit smart phone app	Dairy Australia
2015	Australian Dairy Hygiene Handbook: guidance on high quality raw milk material including management of bacterial contamination	Dairy Australia
2016 (& ongoing)	Dairy Australia Factsheet: Antimicrobial resistance in the dairy industry	https://www.dairyaustralia.com.au/-/media/dairyaustralia/documents/farm/pasture-management/nutrition/advanced-nutrition-in-action/advanced-nutrition-in-action.ashx

2016 (& ongoing)	Dairy Australia “Advanced nutrition in action” promotes understanding of rumen function in transition management and metabolic disease to assist with reduction in use of virginiamycin on a herd level	Dairy Australia
2016 (& ongoing)	Dairy Australia “InCharge workshop” includes a day on transition management and metabolic disease to assist with reduction in use of virginiamycin on a herd level	http://healthyhooves.dairyaustralia.com.au/
2017	Dairy Australia Healthy Hooves support program to manage dairy cow hoof health and reduce reliance on antibiotics	http://drycowconsult.com.au/
2017	Dairy Australia Dry Cow Consult Tool developed to optimize antibiotic use during drying off	http://drycowconsult.com.au/
2017	IDF Factsheet: Guidance on Antimicrobial Resistance from the dairy sector	IDF Factsheet 003/2017-05
2017	Commencement of IDF repository of national strategies and action plans on AMR	Dairy Australia
2017-18	Second survey of antibiotic use in Australian dairy industry by dairy cattle veterinarians (AMU from vet sales data)	Dairy Australia
2018	Development of Biosecurity planning initiative/tool	Dairy Australia

